arXiv:0711.1306v1 [math.NT] 8 Nov 2007

EXPONENTIAL SUMS AND RANK OF PERSYMMETRIC MATRICES OVER $\mathbb{F}_2$

JORGEN CHERLY

Résumé. Soit $\mathbb{K}$ le corps des séries de Laurent formelles $\mathbb{F}_2((T^{-1}))$. Nous calculons en particulier des sommes exponentielles dans $\mathbb{K}$ de la forme $\sum_{degY\leq k-1}\sum_{degZ\leq s-1}E(tYZ)$ où t est dans la boule unité de $\mathbb{K}$, en démontrant qu'elles dépendent seulement du rang de matrices persymétriques avec des entrées dans $\mathbb{F}_2$ qui leur sont associées. (Une matrice $[\alpha_{i,j}]$ est persymétrique si $\alpha_{i,j} = \alpha_{r,s}$ pour i+j = r+s). En outre nous établissons des propriétes de rang d'une partition de matrices persymétriques. Nous utilisons ces résultats pour calculer le nombre Γ_i de matrices persymétriques sur $\mathbb{F}_2$ de rang i. Nous retrouvons en particulier une formule générale donnée par D.E.Daykin. Notre dmonstration est, comme indiqué, très différente, puisqu'elle se fonde sur les propriétés de rang d'une partition de matrices persymétriques. Nous montrons également que le nombre R de représentations dans $\mathbb{F}_2[T]$ de 0 comme une somme de formes quadratiques associées aux sommes exponentielles $\sum_{degY\leq k-1}\sum_{degZ\leq s-1}E(tYZ)$ est donné par une intégrale étendue à la boule unité et est une combinaison linéaire des Γ_i. Nous calculons alors explicitement le nombre R. Des résultats similaires sont également obtenus pour les $\mathbb{K}$- espaces vectoriels de dimension n+1. Nous terminons notre article en calculant explicitement le nombre de matrices de rang i de la forme $\left[\frac{A}{B}\right]$, où A est persymétrique.

The author is indepted to Thierry Levasseur UBO Brest for helpful advice regarding LaTeX.

ABSTRACT. Let $\mathbb{K}$ be the field of Laurent Series $\mathbb{F}_2((T^{-1}))$.
We compute in particular exponential sums in $\mathbb{K}$ of the form
$\sum_{degY\leq k-1}\sum_{degZ\leq s-1}E(tYZ)$ where t is in the unit interval of $\mathbb{K}$, by showing that they only depend on the rank of some associated persymmetric matrices with entries in $\mathbb{F}_2$. (A matrix $[\alpha_{i,j}]$ is persymmetric if $\alpha_{i,j}=\alpha_{r,s}$ for i+j = r+s). Besides we establish rank properties of a partition of persymmetric matrices. We use these results to compute the number Γ_i of persymmetric matrices over $\mathbb{F}_2$ of rank i. We recover in this particular a general formula given by D. E. Daykin. Our proof is as indicated very different since it relies on rank properties of a partition of persymmetric matrices. We also prove that the number R of representations in $\mathbb{F}_2[T]$ of 0 as a sum of some quadratic forms associated to the exponential sums $\sum_{degY\leq k-1}\sum_{degZ\leq s-1}E(tYZ)$ is given by an integral over the unit interval, and is a linear combination of the $\Gamma_i's$. We then compute explicitly the number R. Similar results are also obtained for n+1 dimensional $\mathbb{K}$ - vector spaces.We finish the paper by computing explicitely the number of rank i matrices of the form $\left[\frac{A}{B}\right]$, where A is persymmetric.

Contents

1. Notations 4
1.1. Analysis on $\mathbb{K}$ 4
1.2. Analysis on the (n+1) - dimensional $\mathbb{K}$- vectorspace 6
2. Introduction (refer to Section 1 and Section 3) 8
3. Statement of results 14
4. Exponential sums formulas on $\mathbb{K}$ 22
5. Rank properties of a partition of persymmetric matrices. 26
6. Exponential sums in $\mathbb{K}\times\mathbb{K}$ 31
7. Proof of Theorem 3.1 36
8. Proof of Theorem 3.3 39
9. Proofs of Theorems 3.4,3.5,3.6,3.7 and 3.8 41
9.1. Proof of Theorem 3.4 41
9.2. Proof of Theorem 3.5 41
9.3. Proof of Theorem 3.6 41
9.4. Proof of Theorem 3.7 41
9.5. Proof of Theorem 3.8 41
10. Proofs of Theorem 3.9 , Corollary 3.10 and Theorem 3.11 42
10.1. $\Gamma_{i}^{\left[\substack{n\\1+m}\right]\times k}$ written as a linear combination of the $\Gamma_{i-j}^{(1+m)\times k}$ for $0\leq j\leq n.$ 42
10.2. Computation of a_i^n for $1\leq i\leq n-1$ 44
10.3. Proof of Theorem 3.9 45
10.4. Proof of Corollary 3.10 45
10.5. Proof of Theorem 3.11 45
References 45

1. Notations

1.1. **Analysis on $\mathbb{K}$.** We denote by $\mathbb{F}_2\big(\big(\frac{1}{T}\big)\big) = \mathbb{K}$ the completion of the field $\mathbb{F}_2(T)$, the field of rational fonctions over the finite field $\mathbb{F}_2$, for the infinity valuation $\mathfrak{v} = \mathfrak{v}_\infty$ defined by $\mathfrak{v}\big(\frac{A}{B}\big) = degB - degA$ for each pair (A,B) of non-zero polynomials. Then every element non-zero t in $\mathbb{F}_2\big(\big(\frac{1}{T}\big)\big)$ can be expanded in a unique way in a convergent Laurent series $t = \sum_{j=-\infty}^{-\mathfrak{v}(t)} t_j T^j \ where \ t_j \in \mathbb{F}_2$.
We associate to the infinity valuation $\mathfrak{v} = \mathfrak{v}_\infty$ the absolute value $|\cdot|_\infty$ defined by

$$|t|_\infty = |t| = 2^{-\mathfrak{v}(t)}.$$

We denote E the Character of the additive locally compact group $\mathbb{F}_2\big(\big(\frac{1}{T}\big)\big)$ defined by

$$E\Big(\sum_{j=-\infty}^{-\mathfrak{v}(t)} t_j T^j\Big) = \begin{cases} 1 & \text{if } t_{-1} = 0, \\ -1 & \text{if } t_{-1} = 1. \end{cases}$$

We denote $\mathbb{P}$ the valuation ideal in $\mathbb{K}$, also denoted the unit interval of $\mathbb{K}$, i.e. the open ball of radius 1 about 0 or, alternatively, the set of all Laurent series

$$\sum_{i\geq 1} \alpha_i T^{-i} \quad (\alpha_i \in \mathbb{F}_2)$$

and, for every rational integer j, we denote by $\mathbb{P}_j$ the ideal $\{t \in \mathbb{K} |\ \mathfrak{v}(t) > j\}$. The sets $\mathbb{P}_j$ are compact subgroups of the additive locally compact group $\mathbb{K}$.
All $t \in \mathbb{F}_2\Big(\Big(\frac{1}{T}\Big)\Big)$ may be written in a unique way as $t = [t] + \{t\}$, $[t] \in \mathbb{F}_2[T]$, $\{t\} \in \mathbb{P}(= \mathbb{P}_0)$.
We denote by dt the Haar measure on $\mathbb{K}$ chosen so that

$$\int_{\mathbb{P}} dt = 1.$$

Definition 1.1. We introduce the following definitions in $\mathbb{K}$.

- A matrix $D = [\alpha_{i,j}]$ is said to be persymmetric if $\alpha_{i,j} = \alpha_{r,s}$ whenever i+j = r+s.

- We denote by $Persym_{n\times m}(\mathbb{F}_2)$ the set of all $n \times m$ persymmetric matrices over $\mathbb{F}_2$.

- Let $\Gamma_i^{s\times k}$ denote the number of persymmetric $s \times k$ matrices over $\mathbb{F}_2$ of rank i. We can assume without restriction that $s \leq k$, since obviously the transpose of a persymmetric matrix remains persymmetric.

- Set $t = \sum_{i\geq 1} \alpha_i T^{-i} \in \mathbb{P}$ and let $(l, n, m) \in \mathbb{N}^* \times \mathbb{N}^* \times \mathbb{N}^*$. We denote by $D^l_{n\times m}(t)$ the following $n \times m$ persymmetric matrix

  $$\begin{pmatrix}
  \alpha_l & \alpha_{l+1} & \alpha_{l+2} & \dots & \alpha_{l+m-2} & \alpha_{l+m-1} \\
  \alpha_{l+1} & \alpha_{l+2} & \alpha_{l+3} & \dots & \alpha_{l+m-1} & \alpha_{l+m} \\
  \vdots & \vdots & \vdots & \dots & \vdots & \vdots \\
  \vdots & \vdots & \vdots & \dots & \vdots & \vdots \\
  \alpha_{l+n-2} & \alpha_{l+n-1} & \alpha_{l+n} & \dots & \alpha_{l+n+m-4} & \alpha_{l+n+m-3} \\
  \alpha_{l+n-1} & \alpha_{l+n} & \alpha_{l+n+1} & \dots & \alpha_{l+n+m-3} & \alpha_{l+n+m-2}
  \end{pmatrix}.$$

  If l =1 we denote $D^l_{n\times m}(t)$ by $D_{n\times m}(t)$.

- We denote by ker D the Nullspace of the matrix D and r(D) the rank of the matrix D.

- Let $h_{s,k}(t) = h(t)$ be the quadratic exponential sum defined for rational integers $s, k \geq 1$ by

  $$t \in \mathbb{P} \longmapsto \sum_{degY\leq k-1} \sum_{degZ\leq s-1} E(tYZ) \in \mathbb{Z}.$$

- Let $g_{s,k}(t) = g(t)$ be the quadratic exponential sum defined for rational integers $s, k \geq 2$ by

  $$t \in \mathbb{P} \longmapsto \sum_{degY=k-1} \sum_{degZ=s-1} E(tYZ) \in \mathbb{Z}.$$

- We denote by $\mathbb{P}_{l-1}/\mathbb{P}_{l+n+m-2}$ a complete set of coset representatives of $\mathbb{P}_{l+n+m-2}$ *in* $\mathbb{P}_{l-1}$. Obviously the sums h(t) and g(t) are constant on cosets of $\mathbb{P}_{k+s-1}$.

- Consider the following partition of the matrix $D^l_{n\times m}(t)$, obtained by drawing a horizontal line between the (n-1)th and the nth row and drawing a vertical line between the (m-1)th and the mth column in the matrix $D^l_{n\times m}(t)$.

  $$\left(\begin{array}{ccccc|c}
  \alpha_l & \alpha_{l+1} & \alpha_{l+2} & \dots & \alpha_{l+m-2} & \alpha_{l+m-1} \\
  \alpha_{l+1} & \alpha_{l+2} & \alpha_{l+3} & \dots & \alpha_{l+m-1} & \alpha_{l+m} \\
  \vdots & \vdots & \vdots & \dots & \vdots & \vdots \\
  \vdots & \vdots & \vdots & \dots & \vdots & \vdots \\
  \alpha_{l+n-2} & \alpha_{l+n-1} & \alpha_{l+n} & \dots & \alpha_{l+n+m-4} & \alpha_{l+n+m-3} \\
  \hline
  \alpha_{l+n-1} & \alpha_{l+n} & \alpha_{l+n+1} & \dots & \alpha_{l+n+m-3} & \alpha_{l+n+m-2}
  \end{array}\right).$$

Let$(j_1, j_2, j_3, j_4) \in \mathbb{N}^4$, we define

$$\#\left(\begin{array}{c|c} j_1 & j_2 \\ \hline j_3 & j_4 \end{array}\right)_{\mathbb{P}_{l-1}/\mathbb{P}_{l+n+m-2}}$$

to be the cardinality of the following set

$$\{t \in \mathbb{P}_{l-1}/\mathbb{P}_{l+n+m-2} \mid r(D^{l}_{(n-1)\times(m-1)}(t)) = j_1 \; , r(D^{l}_{(n-1)\times m}(t)) = j_2$$
$$r(D^{l}_{n\times(m-1)}(t)) = j_3 \; , r(D^{l}_{n\times m}(t)) = j_4\}.$$

1.2. **Analysis on the (n+1) - dimensional $\mathbb{K}$- vectorspace.** Let $\mathbb{K}^{n+1}$ be the (n+1)- dimensional vector space over $\mathbb{K}$. Let $(t, \eta_1, \eta_2, \ldots, \eta_n) \in \mathbb{K}^{n+1}$ and $|(t, \eta_1, \eta_2, \ldots, \eta_n)| = sup\,\{|t|, |\eta_1|, |\eta_2|, \ldots, |\eta_n|\} = 2^{-inf(\mathfrak{v}(t),\mathfrak{v}(\eta_1),\mathfrak{v}(\eta_2),\ldots,\mathfrak{v}(\eta_n))}$.

It is easy to see that $(t, \eta_1, \eta_2, \ldots, \eta_n) \longrightarrow |(t, \eta_1, \eta_2, \ldots, \eta_n)|$ is an ultrametric valuation on $\mathbb{K}^{n+1}$, that is

$(t, \eta_1, \eta_2, \ldots, \eta_n) \longrightarrow |(t, \eta_1, \eta_2, \ldots, \eta_n)|$ is a norm and
$|((t, \eta_1, \eta_2, \ldots, \eta_n)+(t', \eta_1', \eta_2', \ldots, \eta_n'))| \leq max\,\{|(t, \eta_1, \eta_2, \ldots, \eta_n)|, |(t', \eta_1', \eta_2', \ldots, \eta_n')|\}$.

We denote by $d((t, \eta_1, \eta_2, \ldots, \eta_n) = dtd\eta_1 d\eta_2 \cdots d\eta_n$ the Haar measure on $\mathbb{K}^{n+1}$ chosen so that the measure on the unit interval of $\mathbb{K}^{n+1}$ is equal to one, that is

$$\int_{\mathbb{P}^{n+1}} d((t, \eta_1, \eta_2, \ldots, \eta_n) = \int_{\mathbb{P}} dt \int_{\mathbb{P}} d\eta_1 \int_{\mathbb{P}} d\eta_2 \cdots \int_{\mathbb{P}} d\eta_n = 1 \cdot 1 \cdot 1 \cdots 1 = 1.$$

Let

$$(t, \eta_1, \eta_2, \ldots, \eta_n) = \Big(\sum_{i=-\infty}^{-\mathfrak{v}(t)} t_i T^i, \sum_{i=-\infty}^{-\mathfrak{v}(\eta_1)} \eta_{1i} T^i, \sum_{i=-\infty}^{-\mathfrak{v}(\eta_2)} \eta_{2i} T^i, \ldots, \sum_{i=-\infty}^{-\mathfrak{v}(\eta_n)} \eta_{ni} T^i \Big) \in \mathbb{K}^{n+1},$$

we denote χ the Character on $(\mathbb{K}^{n+1}, +)$ defined by

$$\chi\Big(\sum_{i=-\infty}^{-\mathfrak{v}(t)} t_i T^i, \sum_{i=-\infty}^{-\mathfrak{v}(\eta_1)} \eta_{1i} T^i, \sum_{i=-\infty}^{-\mathfrak{v}(\eta_2)} \eta_{2i} T^i, \ldots, \sum_{i=-\infty}^{-\mathfrak{v}(\eta_n)} \eta_{ni} T^i \Big)$$
$$= E\Big(\sum_{i=-\infty}^{-\mathfrak{v}(t)} t_i T^i \Big) \cdot E\Big(\sum_{i=-\infty}^{-\mathfrak{v}(\eta_1)} \eta_{1,i} T^i \Big) \cdot E\Big(\sum_{i=-\infty}^{-\mathfrak{v}(\eta_2)} \eta_{2,i} T^i \Big) \cdots E\Big(\sum_{i=-\infty}^{-\mathfrak{v}(\eta_n)} \eta_{n,i} T^i \Big)$$
$$= \begin{cases} 1 & \text{if } t_{-1} + \eta_{1,-1} + \eta_{2,-1} + \ldots + \eta_{n,-1} = 0, \\ -1 & \text{if } t_{-1} + \eta_{1,-1} + \eta_{2,-1} + \ldots + \eta_{n,-1} = 1. \end{cases}$$

Definition 1.2. We introduce the following definitions in the (n+1) - dimensional $\mathbb{K}$- vectorspace.

- We denote by $\mathbb{P}/\mathbb{P}_{k+m}\times(\mathbb{P}/\mathbb{P}_k)^n$ a complete set of coset representatives of $\mathbb{P}_{k+m}\times\mathbb{P}_k^n$ *in* $\mathbb{P}^{n+1}$.

-

$$Set \quad (t,\eta_1,\eta_2,\ldots,\eta_n)=\big(\sum_{i\geq 1}\alpha_i T^i,\sum_{i\geq 1}\beta_{1i}T^i,\sum_{i\geq 1}\beta_{2i}T^i,\ldots,\sum_{i\geq 1}\beta_{ni}T^i\big)\in\mathbb{P}^{n+1}.$$

  We denote by $D^{\left[\substack{n\\1+m}\right]\times k}(t,\eta_1,\eta_2,\ldots,\eta_n)$ the following $(1+n+m)\times k$ matrix, where the submatrix obtained by delating the n last rows form a $(1+m)\times k$ persymmetric matrix, and the n last rows form a $n\times k$ matrix over the finite field $\mathbb{F}_2$

$$\left(\begin{array}{cccccc}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \ldots & \vdots & \vdots \\
\vdots & \vdots & \vdots & \ldots & \vdots & \vdots \\
\alpha_{1+m} & \alpha_{2+m} & \alpha_{3+m} & \ldots & \alpha_{k+m-1} & \alpha_{k+m} \\
\hline
\beta_{11} & \beta_{12} & \beta_{13} & \ldots & \beta_{1k-1} & \beta_{1k} \\
\beta_{21} & \beta_{22} & \beta_{23} & \ldots & \beta_{2k-1} & \beta_{2k} \\
\vdots & \vdots & \vdots & \ldots & \vdots & \vdots \\
\vdots & \vdots & \vdots & \ldots & \vdots & \vdots \\
\beta_{n1} & \beta_{n2} & \beta_{n3} & \ldots & \beta_{nk-1} & \beta_{nk}
\end{array}\right).$$

- Let $g_{m,k}(t,\eta)=g(t,\eta)$ be the exponential sum defined for rational integers $m\geq 0,k\geq 1$ by

$$(t,\eta)\in\mathbb{P}\times\mathbb{P}\longmapsto\sum_{degY\leq k-1}\sum_{degZ\leq m}E(tYZ)\sum_{degU=0}E(\eta YU)\in\mathbb{Z}.$$

- Let $f_{m,k}(t,\eta)=f(t,\eta)$ be the exponential sum defined for rational integers $m\geq 0,k\geq 1$ by

$$(t,\eta)\in\mathbb{P}\times\mathbb{P}\longmapsto\sum_{degY\leq k-1}\sum_{degZ\leq m}E(tYZ)\sum_{degU\leq 0}E(\eta YU)\in\mathbb{Z}.$$

-

$$\begin{aligned}
\sigma_{i,i}^{\left[\substack{1\\1+m}\right]\times k} &= Card\left\{(t,\eta)\in\mathbb{P}/\mathbb{P}_{k+m}\times\mathbb{P}/\mathbb{P}_k\mid r(D_{(1+m)\times k}(t))=r(D^{\left[\substack{1\\1+m}\right]\times k}(t,\eta))=i\right\},\\
\sigma_{i-1,i}^{\left[\substack{1\\1+m}\right]\times k} &= Card\left\{(t,\eta)\in\mathbb{P}/\mathbb{P}_{k+m}\times\mathbb{P}/\mathbb{P}_k\mid r(D_{(1+m)\times k}(t))=i-1,\ r(D^{\left[\substack{1\\1+m}\right]\times k}(t,\eta))=i\right\},\\
\Gamma_i^{\left[\substack{n\\1+m}\right]\times k} &= Card\left\{(t,\eta_1,\eta_2,\ldots,\eta_n)\in\mathbb{P}/\mathbb{P}_{k+m}\times(\mathbb{P}/\mathbb{P}_k)^n\mid r(D^{\left[\substack{n\\1+m}\right]\times k})(t,\eta_1,\eta_2,\ldots,\eta_n)=i\right\}
\end{aligned}$$

2. Introduction (refer to Section 1 and Section 3)

The rational function field $\mathbb{F}_2(T)$ is completed with respect to an appropriate valuation to a field $\mathbb{K}$ (i.e. the field of Laurent Series). The unit interval of $\mathbb{K}$, that is, the open ball of radius 1 about 0, is a compact additive group. We shall use the Haar integral on this group.
The paper is based on the proof of the following formula of the number Γ_i of $s\times k$ persymmetric matrices over $\mathbb{F}_2$ of rank i, that is

$$\Gamma_i^{s\times k} = \left\{ \begin{array}{ccc} 1 & if & i=0, \\ 3\cdot 2^{2(i-1)} & if & 1\leq i\leq s-1, \\ 2^{k+s-1}-2^{2s-2} & if & i=s\ (\leq k). \end{array} \right. \tag{2.1}$$

We remark that David E. Daykin [1] has already proved this result over any finite field $\mathbb{F}$ with the number 2 in the formula replaced by $|\mathbb{F}|$, and the number 3 replaced by $|\mathbb{F}|^2-1$. Our proof is very different since it relies on rank properties of a partition of persymmetric matrices, and is proper to the finite field with two elements. Besides, in order to establish the formula (2.1), we obtain several results concerning exponential sums in $\mathbb{K}$ and persymmetric matrices, which seem to have an intrinsic interest.

The paper is organized as follows:

In Section 1, are introduced main notations and definitions.
In Section 3, we state the main theorems in the (n+1) dimensional $\mathbb{K}$-vector spaces.
In Section 4, we establish in particular results on exponential sums in $\mathbb{P}$ of the form h(t), g(t) and we show that they only depend on rank properties of some corresponding persymmetric matrices with entries in $\mathbb{F}_2$.
The proof of these results is based on the following identity

$$\sum_{degZ\leq s-1} E(tYZ) = \left\{ \begin{array}{ccc} 2^s & if & Y\in \ker D_{s\times k}(t) \\ 0 & & otherwise, \end{array} \right. \tag{2.2}$$

where $t\in\mathbb{P}$ *and* $\deg Y\leq k-1$.
By (2.2) we obtain

(2.3)
$$Let\ t\in\mathbb{P},\quad then\ h_{s,k}(t)=h(t)=\sum_{degY\leq k-1}\sum_{degZ\leq s-1} E(tYZ)\quad is\quad given\quad by\quad 2^{k+s-r(D_{s\times k}(t))}.$$

From (2.3), observing that h(t) is constant on cosets of $\mathbb{P}_{k+s-1}$, we prove that the number R of representations in $\mathbb{F}_2[T]$, of 0 as a sum of quadratic forms is given by an integral over the unit interval of $\mathbb{K}$, and is a linear combination of the $\Gamma_i's$.
More precisely we prove

$$R=\int_{\mathbb{P}} h^q(t)dt = 2^{(q-1)(k+s)+1}\sum_{i=0}^{s}\Gamma_i 2^{-qi}. \tag{2.4}$$

By calculating the square of g(t) and using (2.3) we obtain

(2.5)

$$Let \quad t \in \mathbb{P}, \quad then \quad g_{s,k}(t) = g(t) = \sum_{degY=k-1} \sum_{degZ=s-1} E(tYZ) \quad is \quad given \quad by$$

$$\begin{cases} 2^{s+k-j-2} & \text{if } r(D_{(s-1)\times(k-1)}(t)) = r(D_{s\times(k-1)}(t)) = r(D_{(s-1)\times k}(t)) = r(D_{s\times k}(t)) = j, \\ -2^{s+k-j-2} & \text{if } r(D_{(s-1)\times(k-1)}(t)) = r(D_{s\times(k-1)}(t)) = r(D_{(s-1)\times k}(t)) = j \ and \ r(D_{s\times k}(t)) = j+1, \\ 0 & \text{if } otherwise. \end{cases}$$

In Section 5, we study rank properties of a partition of persymmetric matrices by considering the following partition of the matrix $D_{s\times k}(t)$

$$\left(\begin{array}{ccccc|c} \alpha_1 & \alpha_2 & \alpha_3 & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \alpha_4 & \dots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s-1} & \alpha_s & \alpha_{s+1} & \dots & \alpha_{k+s-3} & \alpha_{k+s-2} \\ \hline \alpha_s & \alpha_{s+1} & \alpha_{s+2} & \dots & \alpha_{k+s-2} & \alpha_{k+s-1} \end{array}\right).$$

We prove

(2.6)

$$If \quad 0 \leq j \leq s-1, \quad then \quad {}^{\#}\left(\begin{array}{c|c} j & j \\ \hline j & j \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} = {}^{\#}\left(\begin{array}{c|c} j & j \\ \hline j & j+1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}.$$

This result is deduced from the fact that

$$\int_{\mathbb{P}} g^{2q+1}(t)dt = 0.$$

$$(2.7) \qquad If \quad 0 \leq j \leq s-2, \quad then \quad {}^{\#}\left(\begin{array}{c|c} j & j+1 \\ \hline j+1 & j+1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} = 0.$$

Actually, this result holds for any matrix.

(2.8)

$$If \quad 0 \leq j \leq s-2, \quad then \quad {}^{\#}\left(\begin{array}{c|c} j & j+1 \\ \hline j & j+1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} = {}^{\#}\left(\begin{array}{c|c} j & j \\ \hline j+1 & j+1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} = 0.$$

These equalities are consequence of rank properties of submatrices of persymmetric $s \times k$ matrices.

By combining (2.6),(2.7) and (2.8) we obtain

(2.9)

$$2\cdot\Gamma_{i}^{s\times(k-1)} = 2\cdot\Gamma_{i}^{(s-1)\times k} = 2\cdot{}^{\#}\left(\begin{array}{c|c} i & i \\ \hline i & i \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} + {}^{\#}\left(\begin{array}{c|c} i-1 & i \\ \hline i & i+1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}.$$

From (2.9) we get

$$(2.10)\qquad If\quad 0\leq i\leq s-2,\quad then\quad \Gamma_{i}^{s\times(k-1)} = \Gamma_{i}^{(s-1)\times k}.$$

In Section 6, we compute exponential sums in $\mathbb{P}^2$ associated to a matrix of the form $\left[\frac{A}{b_-}\right]$, where A is a $(m+1)\times k$ persymmetic matrix over $\mathbb{F}_2$ and b_- a $1\times k$ matrix with entries in $\mathbb{F}_2$.

In particular we prove that the number of rank i matrices of the form

$$\left(\begin{array}{cccccc}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \ldots & \vdots & \vdots \\
\vdots & \vdots & \vdots & \ldots & \vdots & \vdots \\
\alpha_{1+m} & \alpha_{2+m} & \alpha_{3+m} & \ldots & \alpha_{k+m-1} & \alpha_{k+m} \\
\hline
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k
\end{array}\right)$$

is a linear combination of the $\Gamma_{j}^{(1+m)\times k}$ for $j\in\{i-1,i\}$ (recall that $\Gamma_{j}^{(1+m)\times k}$ denotes the number of persymmetric $(1+m)\times k$ matrices of rank j).
In fact we prove that

(2.11)

$$\Gamma_{i}^{\left[\begin{smallmatrix}1\\1+m\end{smallmatrix}\right]\times k} = (2^k-2^{i-1})\cdot\Gamma_{i-1}^{(1+m)\times k}+2^i\Gamma_{i}^{(1+m)\times k}\quad for\quad all\quad 1\leq i\leq inf(k,2+m).$$

The proof of (2.11) is based on the following identity

$$(2.12)\qquad E(\eta Y) = \begin{cases} 1 & \text{if } \sum_{j=1}^{k-1}\beta_j\delta_j = 0 \\ -1 & \text{if } \sum_{j=1}^{k-1}\beta_j\delta_j = 1 \end{cases}$$

where,

$$\eta = \sum_{j\geq 1}\beta_j T^{-j}\in\mathbb{P}\quad and\quad Y = \sum_{i=1}^{k-1}\delta_i T^i\in\mathbb{F}_2[T],\quad degY\leq k-1.$$

We proceed as follows :

By (2.12) we obtain

(2.13)

$$g(t,\eta) = \begin{cases} 2^{k+m+1-r(D_{(1+m)\times k}(t))} & \text{if } r(D_{(1+m)\times k}(t)) = r(D^{\left[\substack{1\\1+m}\right]\times k}(t,\eta)), \\ 0 & \text{otherwise ,} \end{cases}$$

$$where \quad g(t,\eta) = \sum_{degY\leq k-1}\sum_{degZ\leq m} E(tYZ)\sum_{degU=0} E(\eta YU).$$

By squaring $g(t,\eta)$ we obtain $g^2(t,\eta) = g(t,\eta)\cdot h(t)$ where $h(t) = \sum_{deg\leq k-1}\sum_{degZ\leq m} E(tYZ)$. By recurrence on q we get

$$g^q(t,\eta) = g(t,\eta)\cdot h^{q-1}(t). \tag{2.14}$$

By integrating $g^q(t,\eta)$ over the unit interval of $\mathbb{K}^2$, observing that $g(t,\eta)$ is constant on cosets of $\mathbb{P}_{k+m}\times\mathbb{P}_k$ and using (2.13), we obtain

$$\int_{\mathbb{P}}\int_{\mathbb{P}} g^q(t,\eta)dtd\eta = 2^{q(k+m+1)-2k-m}\sum_{i=0}^{inf(k,1+m)}\sigma_{i,i}^{\left[\substack{1\\1+m}\right]\times k}2^{-iq}. \tag{2.15}$$

Integrating $g(t,\eta)\cdot h^{q-1}(t)$ over the unit interval of $\mathbb{K}^2$, we get by Fubini's theorem

$$\int_{\mathbb{P}}\int_{\mathbb{P}} g(t,\eta)h^{q-1}(t)dtd\eta = 2^{q(k+m+1)-2k-m}\sum_{i=0}^{inf(k,1+m)} 2^i\Gamma_i^{(1+m)\times k}2^{-iq}. \tag{2.16}$$

Comparing (2.15) and (2.16), using (2.14), we get

$$\sigma_{i,i}^{\left[\substack{1\\1+m}\right]\times k} = 2^i\Gamma_i^{(1+m)\times k} \quad for\ all\ 0\leq i\leq inf(k,1+m). \tag{2.17}$$

We obviously have

$$\Gamma_i^{\left[\substack{1\\1+m}\right]\times k} = \sigma_{i,i}^{\left[\substack{1\\1+m}\right]\times k} + \sigma_{i-1,i}^{\left[\substack{1\\1+m}\right]\times k}. \tag{2.18}$$

From (2.17) and (2.18) we easily get (2.11).

Besides we prove, similarly to (2.3) and (2.4) that

$$f(t,\eta) = 2^{k+m+2-r(D^{\left[\frac{1}{1+m}\right]\times k}(t,\eta))} \tag{2.19}$$

$$\text{and} \quad \int_{\mathbb{P}}\int_{\mathbb{P}} f^{q}(t,\eta)\,dt\,d\eta = 2^{q(k+m+2)-2k-m}\sum_{i=0}^{inf(k,2+m)}\Gamma_{i}^{\left[\frac{1}{1+m}\right]\times k}2^{-iq} \tag{2.20}$$

$$\text{where} \quad f(t,\eta) = \sum_{degY\leq k-1}\sum_{degZ\leq m}E(tYZ)\sum_{degU\leq 0}E(\eta YU).$$

In Section 7, we prove formula (2.1) by induction on s $(\leq k)$ and by using

$$(2.9) \quad \Gamma_{i}^{s\times(k-1)} = \Gamma_{i}^{(s-1)\times k} \quad for \quad 0\leq i\leq s-2$$

and the two following equations

$$\sum_{i=0}^{s}\Gamma_{i}^{s\times k} = 2^{k+s-1},$$

$$\sum_{i=0}^{s}\Gamma_{i}^{s\times k}\cdot 2^{-i} = 2^{k-1}+2^{s-1}-2^{-1}.$$

In Section 8 we prove Theorem 3.3 by showing that the solutions of the system
(2.21)

$$\Gamma_{i}^{s\times k} = \sum_{(j_1,j_2,j_3)\in\{i-2,i-1,i\}^{3}}\#\left(\begin{array}{c|c} j_1 & j_2 \\ \hline j_3 & i\end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} = \begin{cases} 1 & if \quad i=0, \\ 3\cdot 2^{2(i-1)} & if \quad 1\leq i\leq s-1, \\ 2^{k+s-1}-2^{2s-2} & if \quad i=s \ (\leq k), \end{cases}$$

are given by
(2.22)

$$\#\left(\begin{array}{c|c} j_1 & j_2 \\ \hline j_3 & j_4\end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} = \begin{cases} 1 & \text{if } j_1=j_2=j_3=j_4=0, \\ 2^{2j-1} & \text{if } j_1=j_2=j_3=j\ ,j_4\in\{j,j+1\}\ ,1\leq j\leq s-1, \\ 2^{2j-3} & \text{if } j_1=j-2\ ,j_2=j_3=j-1\ ,j_4=j\ ,2\leq j\leq s, \\ 2^{k+s-1}-2^{2s-1} & \text{if } j_1=j_2=s-1, j_3=j_4=s \\ 0 & \text{otherwise.}\end{cases}$$

In Subsection 9.1, we show that Theorem 3.4 follows from (2.3) and (2.4).

In Subsection 9.2, we show that Theorem 3.5 follows from (2.5) and (2.10).

In Subsection 9.3, we show that Theorem 3.6 follows from (2.13) and (2.15).

In Subsection 9.4, we show that Theorem 3.7 follows from (2.19) and (2.20).

In Subsection 9.5, we show that Theorem 3.8 follows from (2.11) and (2.1) with $s \rightarrow 1+m$.

In Subsection 10.1 , we show that $\Gamma_i^{\left[\substack{n \\ 1+m}\right]\times k}$ can be written as a linear combination of the $\Gamma_{i-j}^{(1+m)\times k}$ for $j = 0,1,\ldots,n,$ where $\Gamma_i^{\left[\substack{n \\ 1+m}\right]\times k}$ denotes the number of rank i matrices of the form $[\frac{A}{B}]$ such that A is a $(1+m)\times k$ persymmetric matrix and B is a $n\times k$ matrix with entries in $\mathbb{F}_2$ and $\Gamma_{i-j}^{(1+m)\times k}$ the number of $(1+m)\times k$ persymmetric matrices of rank i-j.
Precisely we show by induction on n

(2.23)
$$\Gamma_i^{\left[\substack{n \\ 1+m}\right]\times k} = \sum_{j=0}^{n}\Big[2^{(n-j)\cdot(i-j)}a_j^{(n)}\prod_{l=1}^{j}(2^k-2^{i-l})\Big]\cdot\Gamma_{i-j}^{(1+m)\times k} \quad for \quad 0\leq i\leq inf(k,n+m+1)$$

where $a_j^{(n)}$ satisfies the linear recurrence relation

$$a_j^{(n)} = 2^j\cdot a_j^{(n-1)} + a_{j-1}^{(n-1)}, \quad n = 2,3,4,\ldots \quad for \quad 1\leq j\leq n-1.$$

In Subsection 10.2 , the explicit value of $a_j^{(n)}$ in (2.23), is obtained as follows :
Setting m = 0 in (2.23) we have:
(2.24)
$$\Gamma_i^{\left[\substack{n \\ 1}\right]\times k} = \sum_{j=0}^{n}2^{(n-j)\cdot(i-j)}a_j^{(n)}\prod_{l=1}^{j}(2^k-2^{i-l})\Gamma_{i-j}^{1\times k} \quad for \quad 0\leq i\leq inf(k,n+1).$$

Recalling that $\Gamma_i^{\left[\substack{n \\ 1}\right]\times k}$ denotes the number of $(n+1)\times k$ matrices with entries in $\mathbb{F}_2$ of rank i,we get by combining (2.24) and [3, George Landsberg]

$$\Gamma_i^{\left[\substack{n \\ 1}\right]\times k} = \prod_{l=0}^{i-1}\frac{(2^{n+1}-2^l)(2^k-2^l)}{(2^i-2^l)} = \sum_{j=i-1}^{i}2^{(n-j)\cdot(i-j)}a_j^{(n)}\prod_{l=1}^{j}(2^k-2^{i-l})\Gamma_{i-j}^{1\times k}$$

from which, we deduce

$$a_i^{(n)} + 2^{n-(i-1)} \cdot a_{i-1}^{(n)} = \prod_{l=0}^{i-1} \frac{2^{n+1}-2^l}{2^i-2^l}.$$

The formula for $a_i^{(n)}$ given in Lemma 10.2 then follows.
In Subsection 10.3 , we obtain Theorem 3.9 from (2.23) and the explicit value of $a_j^{(n)}$.
In Subsection 10.4 , we get Corollary 3.10 by computing $a_j^{(n)}$ for $0 \leq j \leq n, \quad 1 \leq n \leq 5$.
In Subsection 10.5 , we generalize (2.19) and (2.20).

3. Statement of results

Theorem 3.1. *The number $\Gamma_i^{s\times k}$ of persymmetric $s \times k$ matrices over $\mathbb{F}_2$ of rank i is given by*

$$\Gamma_i^{s\times k} = \begin{cases} 1 & if \quad i = 0, \\ 3 \cdot 2^{2(i-1)} & if \quad 1 \leq i \leq s-1, \\ 2^{k+s-1} - 2^{2s-2} & if \quad i = s \; (s \leq k). \end{cases}$$

Remark 3.2. David E. Daykin has already proved this result over any finite field $\mathbb{F}$ with the number 2 in the formula replaced by$|\mathbb{F}|$, and the number 3 replaced by $|\mathbb{F}|^2 - 1$, see [1]. Our proof is different and proper to the finite field with two elements.

Theorem 3.3. *Let $(j_1, j_2, j_3, j_4) \in \mathbb{N}^4$, then*

$$\#\left(\begin{array}{c|c} j_1 & j_2 \\ \hline j_3 & j_4 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} = \begin{cases} 1 & if \; j_1 = j_2 = j_3 = j_4 = 0, \\ 2^{2j-1} & if \; j_1 = j_2 = j_3 = j \;, j_4 \in \{j, j+1\} \;, 1 \leq j \leq s-1, \\ 2^{2j-3} & if \; j_1 = j-2 \;, j_2 = j_3 = j-1 \;, j_4 = j \;, 2 \leq j \leq s, \\ 2^{k+s-1} - 2^{2s-1} & if \; j_1 = j_2 = s-1, j_3 = j_4 = s, \\ 0 & otherwise, \end{cases}$$

where $\#\left(\begin{array}{c|c} j_1 & j_2 \\ \hline j_3 & j_4 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}$ *was introduced in Definition 1.1.*

Theorem 3.4. *Let $h_{s,k}(t) = h(t)$ be the quadratic exponential sum in $\mathbb{P}$ defined by*

$$t \in \mathbb{P} \longmapsto \sum_{degY \leq k-1} \sum_{degZ \leq s-1} E(tYZ) \in \mathbb{Z}.$$

Then

$$h(t) = 2^{k+s-r(D_{s\times k}(t))}$$

and

$$\int_{\mathbb{P}} h^q(t)dt = 2^{(q-1)(k+s)+1}\sum_{i=0}^{s}\Gamma_i^{s\times k}2^{-qi}.$$

Let R denote the number of solutions $(Y_1, Z_1, \dots, Y_q, Z_q)$ of the polynomial equation

$$Y_1Z_1 + Y_2Z_2 + \ldots + Y_qZ_q = 0$$

satisfying the degree conditions

$$degY_i \leq k-1, \quad degZ_i \leq s-1 \quad for 1 \leq i \leq q.$$

Then

$$R = \int_{\mathbb{P}} h^q(t)dt$$

Theorem 3.5. *Let $g_{s,k}(t) = g(t)$ be the quadratic exponential sum in $\mathbb{P}$ defined by*

$$t \in \mathbb{P} \longmapsto \sum_{degY=k-1}\sum_{degZ=s-1} E(tYZ) \in \mathbb{Z}.$$

Then

$$g(t) = \begin{cases} 2^{s+k-j-2} & if \quad r(D_{(s-1)\times(k-1)}(t)) = r(D_{s\times(k-1)}(t)) = r(D_{(s-1)\times k}(t)) = r(D_{s\times k}(t)) = j, \\ -2^{s+k-j-2} & if \quad r(D_{(s-1)\times(k-1)}(t)) = r(D_{s\times(k-1)}(t)) = r(D_{(s-1)\times k}(t)) = j \text{ and } r(D_{s\times k}(t)) = j+1, \\ 0 & if \quad otherwise, \end{cases}$$

and

$$\int_{\mathbb{P}} g^{2q}(t)dt = 2^{(s+k-2)(2q-1)} \cdot \sum_{j=0}^{s-1} \#\left(\begin{array}{c|c} j & j \\ \hline j & j \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} \cdot 2^{-2qj}.$$

Theorem 3.6 (See Definition 1.2)**.** *Let $g_{m,k}(t,\eta) = g(t,\eta)$ be the exponential sum in $\mathbb{P}\times\mathbb{P}$ defined by*

$$(t,\eta) \in \mathbb{P}\times\mathbb{P} \longmapsto \sum_{degY\leq k-1}\sum_{degZ\leq m} E(tYZ) \sum_{degU=0} E(\eta YU) \in \mathbb{Z}.$$

Then

$$g(t,\eta) = \begin{cases} 2^{k+m+1-r(D_{(1+m)\times k}(t))} & if \, r(D_{(1+m)\times k}(t)) = r(D^{\left[\substack{1\\1+m}\right]\times k}(t,\eta)), \\ 0 & otherwise\ , \end{cases}$$

and

$$\int_{\mathbb{P}}\int_{\mathbb{P}} g^q(t,\eta)dtd\eta = 2^{q(k+m+1)-2k-m}\sum_{i=0}^{inf(k,1+m)} \sigma_{i,i}^{\left[\substack{1\\1+m}\right]\times k}2^{-iq}.$$

Theorem 3.7 (See Definition 1.2)**.** *Let $f_{m,k}(t,\eta) = f(t,\eta)$ be the exponential sum in $\mathbb{P}\times\mathbb{P}$ defined by*

$$(t,\eta)\in\mathbb{P}\times\mathbb{P}\longmapsto\sum_{degY\leq k-1}\sum_{degZ\leq m}E(tYZ)\sum_{degU\leq 0}E(\eta YU)\in\mathbb{Z}.$$

Then

$$f(t,\eta)=2^{k+m+2-r(r(D^{\left[\begin{smallmatrix}1\\1+m\end{smallmatrix}\right]\times k}(t,\eta))}$$

and

$$\int_{\mathbb{P}}\int_{\mathbb{P}}f^{q}(t,\eta)dtd\eta=2^{q(k+m+2)-2k-m}\sum_{i=0}^{inf(k,2+m)}\Gamma_{i}^{\left[\begin{smallmatrix}1\\1+m\end{smallmatrix}\right]\times k}2^{-iq}.$$

Theorem 3.8 (See Definition 1.2)**.** *We have the following formula for all* $0\leq i\leq\inf(k,2+m)$

$$\Gamma_{i}^{\left[\begin{smallmatrix}1\\1+m\end{smallmatrix}\right]\times k}=(2^{k}-2^{i-1})\cdot\Gamma_{i-1}^{(1+m)\times k}+2^{i}\Gamma_{i}^{(1+m)\times k}.$$

The case k = 2

$$\Gamma_{i}^{\left[\begin{smallmatrix}1\\1+m\end{smallmatrix}\right]\times 2}=\begin{cases}1 & \text{if } i=0,\\ 9 & \text{if } i=1,\\ 2^{4+m}-10 & \text{if } i=2.\end{cases}$$

The case m = 0 , k ≥ 2

$$\Gamma_{i}^{\left[\begin{smallmatrix}1\\1\end{smallmatrix}\right]\times k}=\begin{cases}1 & \text{if } i=0,\\ 3\cdot(2^{k}-1) & \text{if } i=1,\\ 2^{2k}-3\cdot 2^{k}+2 & \text{if } i=2.\end{cases}$$

The case m = 1 , k ≥ 3

$$\Gamma_{i}^{\left[\begin{smallmatrix}1\\1+1\end{smallmatrix}\right]\times k}=\begin{cases}1 & \text{if } i=0,\\ 2^{k}+5 & \text{if } i=1,\\ 11\cdot(2^{k}-1) & \text{if } i=2,\\ 2^{2k+1}-3\cdot 2^{k+2}+2^{4} & \text{if } i=3.\end{cases}$$

The case $3 \leq k \leq 1+m$

$$\Gamma_i^{\left[\begin{smallmatrix}1\\1+m\end{smallmatrix}\right]\times k} = \begin{cases} 1 & \textit{if } i=0,\\ 2^k+5 & \textit{if } i=1,\\ 3\cdot 2^{k+2i-4}+21\cdot 2^{3i-5} & \textit{if } 2\leq i\leq k-1,\\ 2^{2k+m}-5\cdot 2^{3k-5} & \textit{if } i=k.\end{cases}$$

The case $2 \leq m \leq k-2$

$$\Gamma_i^{\left[\begin{smallmatrix}1\\1+m\end{smallmatrix}\right]\times k} = \begin{cases} 1 & \textit{if } i=0,\\ 2^k+5 & \textit{if } i=1,\\ 3\cdot 2^{k+2i-4}+21\cdot 2^{3i-5} & \textit{if } 2\leq i\leq m,\\ 11\cdot\left[2^{k+2m-2}-2^{3m-2}\right] & \textit{if } i=m+1,\\ 2^{2k+m}-3\cdot 2^{k+2m}+2^{3m+1} & \textit{if } i=m+2.\end{cases}$$

Theorem 3.9 (See Definitions 1.1, 1.2)**.** *Let* $\Gamma_i^{\left[\begin{smallmatrix}n\\1+m\end{smallmatrix}\right]\times k}$ *denote the number of matrices of the form* $\left[\frac{A}{B}\right]$ *of rank i such that A is a* $(1+m)\times k$ *persymmetric matrix and B is a* $n\times k$ *matrix over* $\mathbb{F}_2$*, and where* $\Gamma_i^{(1+m)\times k}$ *denotes the number of* $(1+m)\times k$ *persymmetric matrices over* $\mathbb{F}_2$ *of rank i.*

Then $\Gamma_i^{\left[\begin{smallmatrix}n\\1+m\end{smallmatrix}\right]\times k}$ *expressed as a linear combination of the* $\Gamma_{i-j}^{(1+m)\times k}$ *is equal to*

$$\sum_{j=0}^{n} 2^{(n-j)\cdot(i-j)} a_j^{(n)} \prod_{l=1}^{j}(2^k-2^{i-l})\cdot\Gamma_{i-j}^{(1+m)\times k} \quad \textit{for} \quad 0\leq i\leq \textit{inf}(k,n+m+1)$$

where

$$a_j^{(n)} = \sum_{s=0}^{j-1}(-1)^s\prod_{l=0}^{j-(s+1)}\frac{2^{n+1}-2^l}{2^{j-s}-2^l}\cdot 2^{s(n-j)+\frac{s(s+1)}{2}}+(-1)^j\cdot 2^{jn-\frac{j(j-1)}{2}} \quad \textit{for} \quad 1\leq j\leq n-1.$$

We set

$$a_0^{(n)} = a_n^{(n)} = 1$$

$$\textit{and} \quad \Gamma_{i-j}^{(1+m)\times k} = 0 \quad \textit{if} \quad i-j\notin\{0,1,2,\ldots,\textit{inf}(k,1+m)\}.$$

Corollary 3.10. *We have the following formulas for n = 1,2,3,4,5 :*

$$\Gamma_i^{\left[\begin{smallmatrix}1\\1+m\end{smallmatrix}\right]\times k} = 2^i\Gamma_i^{(1+m)\times k}+(2^k-2^{i-1})\cdot\Gamma_{i-1}^{(1+m)\times k} \quad \textit{for} \quad 0\leq i\leq \textit{inf}(k,2+m),$$

$$\Gamma_i^{\left[\begin{smallmatrix}2\\1+m\end{smallmatrix}\right]\times k} = 2^{2i}\Gamma_i^{(1+m)\times k}+3\cdot 2^{i-1}(2^k-2^{i-1})\cdot\Gamma_{i-1}^{(1+m)\times k}$$
$$+(2^k-2^{i-1})(2^k-2^{i-2})\cdot\Gamma_{i-2}^{(1+m)\times k} \quad \textit{for} \quad 0\leq i\leq \textit{inf}(k,3+m),$$

$$\Gamma_i^{\left[\begin{smallmatrix}3\\1+m\end{smallmatrix}\right]\times k} = 2^{3i}\Gamma_i^{(1+m)\times k}+7\cdot 2^{(i-1)2}(2^k-2^{i-1})\cdot\Gamma_{i-1}^{(1+m)\times k}$$

$+ 7 \cdot 2^{i-2}(2^k - 2^{i-1})(2^k - 2^{i-2}) \cdot \Gamma_{i-2}^{(1+m)\times k}$
$+ (2^k - 2^{i-1})(2^k - 2^{i-2})(2^k - 2^{i-3})\Gamma_{i-3}^{(1+m)\times k} \quad for \quad 0 \leq i \leq inf(k, 4+m),$

$\Gamma_i^{\left[\begin{smallmatrix}4\\1+m\end{smallmatrix}\right]\times k} = 2^{4i}\Gamma_i^{(1+m)\times k} + 15 \cdot 2^{(i-1)3}(2^k - 2^{i-1}) \cdot \Gamma_{i-1}^{(1+m)\times k}$
$+ 35 \cdot 2^{2i-4}(2^k - 2^{i-1})(2^k - 2^{i-2}) \cdot \Gamma_{i-2}^{(1+m)\times k}$
$+ 15 \cdot 2^{i-3}(2^k - 2^{i-1})(2^k - 2^{i-2})(2^k - 2^{i-3})\Gamma_{i-3}^{(1+m)\times k}$
$+ (2^k - 2^{i-1})(2^k - 2^{i-2})(2^k - 2^{i-3})(2^k - 2^{i-4})\Gamma_{i-4}^{(1+m)\times k}$
$for \quad 0 \leq i \leq inf(k, 5+m),$

$\Gamma_i^{\left[\begin{smallmatrix}5\\1+m\end{smallmatrix}\right]\times k} = 2^{5i}\Gamma_i^{(1+m)\times k} + 31 \cdot 2^{(i-1)4}(2^k - 2^{i-1}) \cdot \Gamma_{i-1}^{(1+m)\times k}$
$+ 155 \cdot 2^{3i-6}(2^k - 2^{i-1})(2^k - 2^{i-2}) \cdot \Gamma_{i-2}^{(1+m)\times k}$
$+ 155 \cdot 2^{2i-6}(2^k - 2^{i-1})(2^k - 2^{i-2})(2^k - 2^{i-3})\Gamma_{i-3}^{(1+m)\times k}$
$+ 31 \cdot 2^{i-4}(2^k - 2^{i-1})(2^k - 2^{i-2})(2^k - 2^{i-3})(2^k - 2^{i-4})\Gamma_{i-4}^{(1+m)\times k}$
$+ (2^k - 2^{i-1})(2^k - 2^{i-2})(2^k - 2^{i-3})(2^k - 2^{i-4})(2^k - 2^{i-5})\Gamma_{i-5}^{(1+m)\times k}$
$for \quad 0 \leq i \leq inf(k, 6+m).$

Theorem 3.11. *(See Section 1) Let* $f_{m,k}(t, \eta_1, \eta_2, \dots, \eta_n)$ *be the exponential sum in* $\mathbb{P}^{n+1}$ *defined by*
$(t, \eta_1, \eta_2, \dots, \eta_n) \in \mathbb{P}^{n+1} \longrightarrow$
$\sum_{degY \leq k-1} \sum_{degZ \leq m} E(tYZ) \sum_{degU_1 \leq 0} E(\eta_1 YU_1) \sum_{degU_2 \leq 0} E(\eta_2 YU_2) \dots \sum_{degU_n \leq 0} E(\eta_n YU_n).$

Set

$$(t, \eta_1, \eta_2, \dots, \eta_n) = \big(\sum_{i \geq 1} \alpha_i T^{-i}, \sum_{i \geq 1} \beta_{1i} T^{-i}, \dots, \sum_{i \geq 1} \beta_{ni} T^{-i}\big) \in \mathbb{P}^{n+1}.$$

Then

$$f_{m,k}(t, \eta_1, \eta_2, \dots, \eta_n) = 2^{k+m+n+1-r(D^{\left[\begin{smallmatrix}n\\1+m\end{smallmatrix}\right]\times k}(t,\eta_1,\eta_2,\dots,\eta_n))}$$

where

$$D^{\left[\begin{smallmatrix}n\\1+m\end{smallmatrix}\right]\times k}(t, \eta_1, \eta_2, \dots, \eta_n)$$

denotes the following $(1+n+m)\times k$ *matrix*

$$\left(\begin{array}{cccccc}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \ldots & \vdots & \vdots \\
\vdots & \vdots & \vdots & \ldots & \vdots & \vdots \\
\alpha_{1+m} & \alpha_{2+m} & \alpha_{3+m} & \ldots & \alpha_{k+m-1} & \alpha_{k+m} \\
\hline
\beta_{11} & \beta_{12} & \beta_{13} & \ldots & \beta_{1k-1} & \beta_{1k} \\
\beta_{21} & \beta_{22} & \beta_{23} & \ldots & \beta_{2k-1} & \beta_{2k} \\
\vdots & \vdots & \vdots & \ldots & \vdots & \vdots \\
\vdots & \vdots & \vdots & \ldots & \vdots & \vdots \\
\beta_{n1} & \beta_{n2} & \beta_{n3} & \ldots & \beta_{nk-1} & \beta_{nk}
\end{array}\right).$$

Then the number denoted by $R_q(n,k,m)$ *of solutions*

$$(Y_1,Z_1,U_1^{(1)},U_2^{(1)},\ldots,U_n^{(1)},Y_2,Z_2,U_1^{(2)},U_2^{(2)},\ldots,U_n^{(2)},\ldots Y_q,Z_q,U_1^{(q)},U_2^{(q)},\ldots,U_n^{(q)})$$

of the polynomial equations

$$\begin{cases}
Y_1Z_1+Y_2Z_2+\ldots+Y_qZ_q=0\\
Y_1U_1^{(1)}+Y_2U_1^{(2)}+\ldots+Y_qU_1^{(q)}=0\\
Y_1U_2^{(1)}+Y_2U_2^{(2)}+\ldots+Y_qU_2^{(q)}=0\\
\vdots\\
Y_1U_n^{(1)}+Y_2U_n^{(2)}+\ldots+Y_qU_n^{(q)}=0
\end{cases}$$

satisfying the degree conditions

$$degY_i\leq k-1,\quad degZ_i\leq m,\quad degU_j^i\leq 0,\quad for\quad 1\leq j\leq n\quad 1\leq i\leq q$$

is equal to the following integral over the unit interval in $\mathbb{K}^{n+1}$

$$\int_{\mathbb{P}^{n+1}} f_{m,k}^q(t,\eta_1,\eta_2,\ldots,\eta_n)\,dt\,d\eta_1 d\eta_2\ldots d\eta_n.$$

Observing that $f_{m,k}(t,\eta_1,\eta_2,\ldots,\eta_n)$ *is constant on cosets of* $\mathbb{P}_{k+m}\times\mathbb{P}_k^n$, *the above integral is equal to*

$$2^{q(k+m+n+1)-(n+1)k-m}\sum_{i=0}^{inf(k,n+1+m)}\Gamma_i^{\left[\substack{n\\1+m}\right]\times k}2^{-iq}=R_q(n,k,m)$$

Example. The number $R_q(0,k,m)$ of solutions $(Y_1,Z_1,\ldots,Y_q,Z_q)$ of the polynomial equation

$$Y_1Z_1+Y_2Z_2+\ldots+Y_qZ_q=0$$

satisfying the degree conditions

$$degY_i \leq k-1, \quad degZ_i \leq m \leq k-1 \quad for 1 \leq i \leq q.$$

is equal to the following integral

$$\int_{\mathbb{P}} \Big[\sum_{degY \leq k-1} \sum_{degZ \leq m} E(tYZ)\Big]^q dt = 2^{(q-1)(k+m+1)+1} \sum_{i=0}^{1+m} \Gamma_i^{(1+m)\times k} 2^{-qi}$$

$$= \begin{cases} 2^k + 2^{1+m} - 1 & if \quad q=1, \\ 2^{2k} + 3\cdot(m+1)\cdot 2^{k+m} & if \quad q=2, \\ 2^{(q-1)(k+m+1)+1}\left[1 + 3\frac{1-2^{(2-q)m}}{2^q-2^2} + (2^{k+m} - 2^{2m})2^{-q(1+m)}\right] & if \quad 3 \leq q. \end{cases}$$

Example. The number $\Gamma_i^{\left[\substack{1 \\ 1+2}\right]\times 3}$ of rank i matrices of the form

$$\left(\begin{array}{ccc} \alpha_1 & \alpha_2 & \alpha_3 \\ \alpha_2 & \alpha_3 & \alpha_4 \\ \alpha_3 & \alpha_4 & \alpha_5 \\ \hline \beta_1 & \beta_2 & \beta_3 \end{array}\right)$$

is equal to

$$\begin{cases} 1 & \text{if } i=0, \\ 13 & \text{if } i=1, \\ 66 & \text{if } i=2, \\ 176 & \text{if } i=3. \end{cases}$$

The number $R_q(1,3,2)$ of solutions $(Y_1, Z_1, U_1, \ldots, Y_q, Z_q, U_q)$ of the polynomial equations

$$\begin{cases} Y_1Z_1 + Y_2Z_2 + \ldots + Y_qZ_q = 0 \\ Y_1U_1 + Y_2U_2 + \ldots + Y_qU_q = 0 \end{cases}$$

satisfying the degree conditions

$$degY_i \leq 2, \quad degZ_i \leq 2, \quad degU_i \leq 0 \quad for \quad 1 \leq i \leq q$$

is equal to the following integral

$$\int_{\mathbb{P}}\int_{\mathbb{P}} \big[\sum_{degY \leq 2} \sum_{degZ \leq 2} E(tYZ) \sum_{degU \leq 0} E(\eta YU)\big]^q dt d\eta = 2^{7q-8} \sum_{i=0}^{3} \Gamma_i^{\left[\substack{1 \\ 1+2}\right]\times 3} 2^{-iq}$$

$$= 2^{4q-8} \cdot \big[2^{3q} + 13\cdot 2^{2q} + 66\cdot 2^q + 176\big].$$

Example. The number $\Gamma_i^{\left[\substack{5\\1+2}\right]\times 4}$ of rank i matrices of the form

$$\left(\begin{array}{cccc}
\alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 \\
\alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 \\
\alpha_3 & \alpha_4 & \alpha_5 & \alpha_6 \\
\hline
\beta_{11} & \beta_{12} & \beta_{13} & \beta_{14} \\
\beta_{21} & \beta_{22} & \beta_{23} & \beta_{24} \\
\beta_{31} & \beta_{32} & \beta_{33} & \beta_{34} \\
\beta_{41} & \beta_{42} & \beta_{43} & \beta_{44} \\
\beta_{51} & \beta_{52} & \beta_{53} & \beta_{54}
\end{array}\right)$$

is equal to

$$\begin{cases}
1 & \text{if } i = 0, \\
561 & \text{if } i = 1, \\
65670 & \text{if } i = 2, \\
3731208 & \text{if } i = 3, \\
63311424 & \text{if } i = 4.
\end{cases}$$

The number $R_3(5,4,2)$ of solutions

$$(Y_1, Z_1, U_1^{(1)}, U_2^{(1)}, U_3^{(1)}, U_4^{(1)}, U_5^{(1)}, Y_2, Z_2, U_1^{(2)}, U_2^{(2)}, U_3^{(2)}, U_4^{(2)}, U_5^{(2)}, Y_3, Z_3, U_1^{(3)}, U_2^{(3)}, U_3^{(3)}, U_4^{(3)}, U_5^{(3)})$$

of the polynomial equations

$$\begin{cases}
Y_1Z_1 + Y_2Z_2 + Y_3Z_3 = 0, \\
Y_1U_1^{(1)} + Y_2U_1^{(2)} + Y_3U_1^{(3)} = 0, \\
Y_1U_2^{(1)} + Y_2U_2^{(2)} + Y_3U_2^{(3)} = 0, \\
Y_1U_3^{(1)} + Y_2U_3^{(2)} + Y_3U_3^{(3)} = 0, \\
Y_1U_4^{(1)} + Y_2U_4^{(2)} + Y_3U_4^{(3)} = 0, \\
Y_1U_5^{(1)} + Y_2U_5^{(2)} + Y_3U_5^{(3)} = 0,
\end{cases}$$

satisfying the degree conditions

$$degY_i \leq 3, \quad degZ_i \leq 2, \quad degU_i \leq 0 \quad for \quad 1 \leq j \leq 5 \quad 1 \leq i \leq 3$$

is equal to the following integral over the unit interval in $\mathbb{K}^6$

$$\int_{\mathbb{P}^6} f_{2,4}^3(t, \eta_1, \eta_2, \eta_3, \eta_4, \eta_5)dtd\eta_1 d\eta_2 d\eta_3 d\eta_4 d\eta_5 = 2^{10} \cdot \sum_{i=0}^{4} \Gamma_i^{\left[\substack{5\\1+2}\right]\times 4} 2^{-i3} = 24413824.$$

4. EXPONENTIAL SUMS FORMULAS ON $\mathbb{K}$

In this section we compute exponential quadratic sums in $\mathbb{P}$ and show that they only depend on rank properties of some associated persymmetric matrices. The following propositions are proved in [2].

Proposition 4.1. *The following holds*

- *For every rational integer j , the measure of $\mathbb{P}_j$ is 2^{-j}.*
- *For every $A \in \mathbb{F}_2[T]$, $E(A) = 1$.*
- *For $u \in \mathbb{K}$ $\nu(u) \geq 2 \Rightarrow$ $E(u) = 1$.*

Proposition 4.2. *Let j be a rational integer and $u \in \mathbb{K}$, then*

$$\int_{\mathbb{P}_j} E(ut)dt = \begin{cases} 2^{-j} & if \quad \nu(u) > -j, \\ 0 & otherwise. \end{cases}$$

Proposition 4.3. *Let j be a rational integer and let $u \in \mathbb{K}$, then*

$$\sum_{degB \leq j} E(Bu) = \begin{cases} 2^{j+1} & if \quad \nu(\{u\}) > j+1, \\ 0 & otherwise. \end{cases}$$

Lemma 4.4. *Let $Y \in \mathbb{F}_2[T]$, then*

$$\sum_{degZ \leq s-1} E(tYZ) = \begin{cases} 2^{s} & if \quad \nu(\{tY\}) > s, \\ 0 & otherwise. \end{cases}$$

Proof. Lemma 4.4 follows from Proposition 4.3 with u = tY and j = s -1. ☐

Lemma 4.5. *Let $t \in \mathbb{P}$ and $Y \in \mathbb{F}_2[T]$, $degY \leq k-1$, then*

$$\nu(\{tY\}) > s \Longleftrightarrow Y \in \ker D_{s\times k}(t).$$

Proof. Let $Y = \sum_{j=0}^{k-1} \gamma_j T^j$, $\gamma_j \in \mathbb{F}_2$, then

$$tY = (\sum_{i\geq 1} \alpha_i T^{-i})(\sum_{j=0}^{k-1} \gamma_j T^j) = \sum_{i\geq 1}\sum_{j=0}^{k-1} \alpha_i \gamma_j T^{-(i-j)}$$

and

$$\{tY\} = \left(\sum_{i=1}^{k} \alpha_i\gamma_{i-1}\right)T^{-1} + \left(\sum_{i=2}^{k+1} \alpha_i\gamma_{i-2}\right)T^{-2} + \ldots + \left(\sum_{i=s}^{k+s-1} \alpha_i\gamma_{i-s}\right)T^{-s} + \ldots$$

Therefore $\nu(\{tY\}) \geq s$ if and only if

$$\begin{bmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 & \ldots & \alpha_k \\ \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 & \alpha_6 & \ldots & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ldots & \vdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ldots & \vdots \\ \alpha_s & \alpha_{s+1} & \alpha_{s+2} & \alpha_{s+3} & \alpha_{s+4} & \ldots & \alpha_{k+s-1} \end{bmatrix} \begin{bmatrix} \gamma_0 \\ \gamma_1 \\ \gamma_2 \\ \gamma_3 \\ \gamma_4 \\ \vdots \\ \gamma_{k-1} \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

$$\Longleftrightarrow D_{s\times k}(t)\begin{pmatrix}\gamma_0\\ \gamma_1\\ \vdots\\ \gamma_{k-1}\end{pmatrix}=\begin{pmatrix}0\\0\\ \vdots\\ 0\end{pmatrix}\Longleftrightarrow Y=\begin{pmatrix}\gamma_0\\ \gamma_1\\ \vdots\\ \gamma_{k-1}\end{pmatrix}\in Ker\,D_{s\times k}(t).$$

□

Lemma 4.6. *Let $t\in\mathbb{P}$ and let q be a rational integer ≥ 1, then*

$$h_{s,k}(t)=h(t)=\sum_{degY\leq k-1}\sum_{degZ\leq s-1}E(tYZ)=2^{k+s-r\left(D_{s\times k}(t)\right)}$$

and

$$\int_{\mathbb{P}}h^q(t)=2^{(q-1)\cdot(k+s)+1}\sum_{i=0}^{s}\Gamma_i^{s\times k}2^{-qi}.$$

Proof. By Lemma 4.4 and Lemma 4.5 we obtain

$$\sum_{degY\leq k-1}\sum_{degZ\leq s-1}E(tYZ)=2^s\sum_{\substack{degY\leq k-1\\ Y\in Ker D_{s\times k}(t)}}1=2^s\cdot 2^{k-r\left(D_{s\times k}(t)\right)}=2^{s+k-r\left(D_{s\times k}(t)\right)}.$$

Observing that h(t) is constant on cosets of $\mathbb{P}_{k+s-1}$, we obtain by integrating $h^q(t)$ on the unit interval of $\mathbb{K}$

$$\begin{aligned}\int_{\mathbb{P}}h^q(t)dt&=\sum_{t\in\mathbb{P}/\mathbb{P}_{k+s-1}}2^{[s+k-r(D_{s\times k}(t))]\cdot q}\int_{\mathbb{P}_{k+s-1}}dt\\&=\sum_{i=0}^{s}\Gamma_i^{s\times k}\cdot 2^{(s+k-i)\cdot q}\int_{\mathbb{P}_{k+s-1}}dt\\&=\sum_{i=0}^{s}\Gamma_i^{s\times k}\cdot 2^{(s+k-i)\cdot q}\cdot 2^{-(k+s-1)}\\&=2^{(q-1)(k+s)+1}\sum_{i=0}^{s}\Gamma_i 2^{-qi}.\end{aligned}$$

□

Lemma 4.7. *Let $t\in\mathbb{P}$ and $g_{s,k}(t)=g(t)=\sum_{degY=k-1}\sum_{degZ=s-1}E(tYZ)$, then*

$$g^2(t)=g_1(t)\cdot g_2(t)\quad where$$

$g_1(t)=\sum_{degY\leq k-2}\sum_{degZ=s-1}E(tYZ)$ *and* $\quad g_2(t)=\sum_{degY=k-1}\sum_{degZ\leq s-2}E(tYZ).$

Proof. Observe that

$$g^2(t) = \sum_{degY=k-1} \sum_{degY_1=k-1} \sum_{degZ=s-1} \sum_{degZ_1=s-1} E(tYZ)E(tY_1Z_1).$$

Set

$$\begin{cases} Y+Y_1 = Y_2 & degY_2 \leq k-2, \\ Y_1 = Y_3 & degY_3 = k-1, \\ Z+Z_1 = Z_2 & degZ_2 \leq s-2, \\ Z_1 = Z_3 & degZ_3 = s-1. \end{cases}$$

Then we obtain

$$\begin{aligned} g^2(t) &= \sum_{degY_2\leq k-2} \sum_{degY_3=k-1} \sum_{degZ_2\leq s-2} \sum_{degZ_3=s-1} E(t(Y_2+Y_3)Z_3)E(tY_3(Z_2+Z_3)) \\ &= \sum_{degY_2\leq k-2} \sum_{degY_3=k-1} \sum_{degZ_2\leq s-2} \sum_{degZ_3=s-1} E(tY_2Z_3)E(tY_3Z_2) \\ &= \Big[\sum_{degY\leq k-2} \sum_{degZ=s-1} E(tYZ)\Big] \cdot \Big[\sum_{degY=k-1} \sum_{degZ\leq s-2} E(tYZ)\Big] \\ &= g_1(t) \cdot g_2(t). \end{aligned}$$

□

Lemma 4.8. *Let* $t \in \mathbb{P}$, *then*

$$g_1(t) = \begin{cases} 2^{k+s-2-r(D_{(s-1)\times(k-1)}(t))} & \text{if } r(D_{(s-1)\times(k-1)}(t)) = r(D_{s\times(k-1)}(t)), \\ 0 & \text{otherwise }, \end{cases}$$

and

$$g_2(t) = \begin{cases} 2^{k+s-2-r(D_{(s-1)\times(k-1)}(t))} & \text{if } r(D_{(s-1)\times(k-1)}(t)) = r(D_{(s-1)\times k}(t)), \\ 0 & \text{otherwise }. \end{cases}$$

Proof. Observe that

$$\begin{aligned} g_1(t) &= \sum_{degY\leq k-2} \sum_{degZ=s-1} E(tYZ) \\ &= \sum_{degY\leq k-2} \sum_{degZ\leq s-1} E(tYZ) - \sum_{degY\leq k-2} \sum_{degZ\leq s-2} E(tYZ) \\ &= h_{s,k-1}(t) - h_{s-1,k-1}(t). \end{aligned}$$

By Lemma 4.6 we obtain

$$g_1(t) = 2^{k+s-1-r\left(D_{s\times(k-1)}(t)\right)} - 2^{k+s-2-r\left(D_{(s-1)\times(k-1)}(t)\right)}.$$

In the same way we obtain

$$g_2(t) = 2^{k+s-1-r\left(D_{(s-1)\times k}(t)\right)} - 2^{k+s-2-r\left(D_{(s-1)\times(k-1)}(t)\right)}$$

and Lemma 4.8 is proved. ☐

Lemma 4.9. *We have the following equivalences*

$$\begin{aligned} g(t) \neq 0 &\iff g^2(t) \neq 0 \\ &\iff g_1(t) \cdot g_2(t) \neq 0 \\ &\iff r\big(D_{(s-1)\times(k-1)}(t)\big) = r\big(D_{s\times(k-1)}(t)\big) = r\big(D_{(s-1)\times k}(t)\big) \end{aligned}$$

Proof. Lemma 4.9 is obtained by combining Lemma 4.7 and Lemma 4.8 ☐

Lemma 4.10. *Let $t \in \mathbb{P}$, then $g_{s,k}(t) = g(t)$ is given by*

- $2^{s+k-j-2}$ *if* $r(D_{(s-1)\times(k-1)}(t)) = r(D_{s\times(k-1)}(t)) = r(D_{(s-1)\times k}(t)) = r(D_{s\times k}(t)) = j$
- $-2^{s+k-j-2}$ *if* $r(D_{(s-1)\times(k-1)}(t)) = r(D_{s\times(k-1)}(t)) = r(D_{(s-1)\times k}(t)) = j$ *and* $r(D_{s\times k}(t)) = j+1$
- *0 otherwise.*

Proof. We see that

$$g(t) = \sum_{degY=k-1}\sum_{degZ=s-1} E(tYZ) = \sum_{degY\leq k-1}\sum_{degZ=s-1} E(tYZ) - \sum_{degY\leq k-2}\sum_{degZ=s-1} E(tYZ).$$

By Lemma 4.9

$$g(t) \neq 0 \iff r\big(D_{(s-1)\times(k-1)}(t)\big) = r\big(D_{s\times(k-1)}(t)\big) = r\big(D_{(s-1)\times k}(t)\big).$$

We consider the following two cases:
First case

$$r\big(D_{(s-1)\times(k-1)}(t)\big) = r\big(D_{s\times(k-1)}(t)\big) = r\big(D_{(s-1)\times k}(t)\big) = r\big(D_{s\times k}(t)\big) = j.$$

Second case

$$r\big(D_{(s-1)\times(k-1)}(t)\big) = r\big(D_{s\times(k-1)}(t)\big) = r\big(D_{(s-1)\times k}(t)\big) = j$$

and

$$r\big(D_{s\times k}(t)\big) = j+1.$$

By the results of lemma 4.8 we obtain:
In the first case

$$\sum_{degY\leq k-1}\sum_{degZ=s-1} E(tYZ) - \sum_{degY\leq k-2}\sum_{degZ=s-1} E(tYZ)$$

$$= 2^{k+s-1-j} - 2^{k+s-2-j} = 2^{k+s-2-j}.$$

In the second case

$$\sum_{degY\leq k-1}\sum_{degZ=s-1} E(tYZ) - \sum_{degY\leq k-2}\sum_{degZ=s-1} E(tYZ)$$

$$= 0 - 2^{k+s-2-j} = -2^{k+s-2-j}$$

and Lemma 4.10 is proved. $\square$

5. RANK PROPERTIES OF A PARTITION OF PERSYMMETRIC MATRICES.

[See Definition 1.1]. In this section we establish rank properties of a partition of the following $s \times k$ persymmetric matrix, obtained by drawing in the matrix a horizontal line between the (s -1)th and the sth row and drawing a vertical line between the (k-1)th and the kth column

$$t = \sum_{i \geq 1} \alpha_i T^{-1} \in \mathbb{P} \longrightarrow \left(\begin{array}{cccccc|c} \alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & & \alpha_k \\ \alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & & \vdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & & \vdots \\ \alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{k+s-3} & & \alpha_{k+s-2} \\ \hline \alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{k+s-2} & & \alpha_{k+s-1} \end{array} \right) \in Persym_{s \times k}(\mathbb{F}_2)$$

Lemma 5.1. *Let* $q \in \mathbb{N}$, *then*

$$\int_{\mathbb{P}} g^{2q+1}(t)dt = 0.$$

Proof. Recall that by Lemma 4.7

$$g_{s,k}(t) = g(t) = \sum_{degY=k-1} \sum_{degZ=s-1} E(tYZ),$$

then

$$\int_{\mathbb{P}} g^{2q+1}(t)dt$$

$$= \sum_{degY_1=k-1} \sum_{degZ_1=s-1} \cdots \sum_{degY_{2q+1}=k-1} \sum_{degZ_{2q+1}=s-1} \int_{\mathbb{P}} \prod_{i=1}^{2q+1} E(tY_iZ_i)dt$$

$$= \sum_{degY_1=k-1} \sum_{degZ_1=s-1} \cdots \sum_{degY_{2q+1}=k-1} \sum_{degZ_{2q+1}=s-1} \int_{\mathbb{P}} E\Big(t \sum_{i=1}^{2q+1} Y_iZ_i\Big)dt.$$

By Proposition 4.2 with j = 0

$$\int_{\mathbb{P}} E(t(Y_1Z_1 + Y_2Z_2 + \ldots + Y_{2q+1}Z_{2q+1}))dt = 0 \Longleftrightarrow \sum_{i=1}^{2q+1} Y_iZ_i = 0$$

but 2q +1 is odd, therefore $deg\Big(\sum_{i=1}^{2q+1} Y_iZ_i \Big) = s + k - 2.$ $\square$

Lemma 5.2. *Let $j \in \mathbb{N}$ such that $0 \leq j \leq s-1$, then*

$$\#\left(\begin{array}{c|c} j & j \\ \hline j & j \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} = \#\left(\begin{array}{c|c} j & j \\ \hline j & j+1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}$$

(see Definition 1.1).

Proof. We define

$\mathbb{A} = \{t \in \mathbb{P} \mid r(D_{(s-1)\times(k-1)}(t)) = r(D_{(s-1)\times k}(t)) = r(D_{s\times(k-1)}(t)) = r(D_{s\times k}(t))\}$

and

$\mathbb{B} = \{t \in \mathbb{P} \mid r(D_{(s-1)\times(k-1)}(t)) = r(D_{(s-1)\times k}(t)) = r(D_{s\times(k-1)}(t)), r(D_{s\times k}(t)) = r(D_{(s-1)\times(k-1)}(t))+1\}.$

By lemma 4.10 we have by observing that g(t) is constant on cosets of $\mathbb{P}_{k+s-1}$

$$\int_{\mathbb{P}} g^{2q+1}(t)dt = \int_{\mathbb{A}} g^{2q+1}(t)dt + \int_{\mathbb{B}} g^{2q+1}(t)dt$$

$$= \sum_{t\in\mathbb{A}\bigcap\left(\mathbb{P}/\mathbb{P}_{k+s-1}\right)} 2^{(s+k-2-r(D_{(s-1)\times(k-1)}(t)))(2q-1)} \int_{\mathbb{P}_{s+k-1}} dt$$

$$+ \sum_{t\in\mathbb{B}\bigcap\left(\mathbb{P}/\mathbb{P}_{k+s-1}\right)} -2^{(s+k-2-r(D_{(s-1)\times(k-1)}(t)))(2q+1)} \int_{\mathbb{P}_{s+k-1}} dt$$

$$= \sum_{j=0}^{s-1} 2^{(s+k-2-j)(2q+1)} \cdot \#\left(\begin{array}{c|c} j & j \\ \hline j & j \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} \int_{\mathbb{P}_{k+s-1}} dt$$

$$-\sum_{j=0}^{s-1} 2^{(s+k-2-j)(2q+1)} \cdot \#\left(\begin{array}{c|c} j & j \\ \hline j & j+1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} \int_{\mathbb{P}_{k+s-1}} dt$$

By lemma 5.1, for all $q \in \mathbb{N}$

$$\sum_{j=0}^{s-1} 2^{(-j)(2q+1)} \cdot \left(\#\left(\begin{array}{c|c} j & j \\ \hline j & j \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} - \#\left(\begin{array}{c|c} j & j \\ \hline j & j+1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}\right) = 0.$$

This completes the proof. ☐

Lemma 5.3 (See Definition 1.1)**.** *Let $j \in \mathbb{N}$ such that $0 \leq j \leq s-2$, then*

$$\#\left(\begin{array}{c|c} j & j+1 \\ \hline j+1 & j+1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} = 0, \tag{5.1}$$

(5.2)

$$\left.\begin{array}{c} r(D_{(s-1)\times(k-1)}(t)) = r(D_{s\times(k-1)}(t)) = j \\ r(D^2_{(s-1)\times(k-2)}(t)) = j-1 \end{array}\right\} \Longrightarrow r(D^2_{s\times(k-2)}(t)) = j-1.$$

Proof. We consider the following partition of the matrix $D_{s\times k}(t)$

$$\left(\begin{array}{c|cccc|c}
\alpha_1 & \alpha_2 & \alpha_3 & \dots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \dots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \dots & \alpha_{k+s-3} & \alpha_{k+s-2} \\
\hline
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \dots & \alpha_{k+s-2} & \alpha_{k+s-1}
\end{array}\right)$$

Obviously we have $r(D_{(s-1)\times k}(t)) = r(D_{s\times k}(t)) \Longrightarrow r(D_{(s-1)\times(k-1)}(t)) = r(D_{s\times(k-1)}(t))$ and so (5.1) is proved. We remark that this result is true for any matrix. To prove (5.2) we consider the matrix obtained by the matrix $D_{s\times k}(t)$ by deleting the first column and replacing the last column by the first one :

$$\left(\begin{array}{cccc|c}
\alpha_2 & \alpha_3 & \dots & \alpha_{k-1} & \alpha_1 \\
\alpha_3 & \alpha_4 & \dots & \alpha_k & \alpha_2 \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_s & \alpha_{s+1} & \dots & \alpha_{k+s-3} & \alpha_{s-1} \\
\hline
\alpha_{s+1} & \alpha_{s+2} & \dots & \alpha_{k+s-2} & \alpha_s
\end{array}\right).$$

Now (5.2) follows from (5.1) with j replaced by j-1. $\square$

Lemma 5.4. *For every $j \in \mathbb{N}$ such that $0 \leq j \leq s-2$ we have*

$$\#\left(\begin{array}{c|c} j & j+1 \\ \hline j & j+1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} = \#\left(\begin{array}{c|c} j & j \\ \hline j+1 & j+1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} = 0$$

(see Definition 1.1).

Proof. Proof by contradiction.
Assume on the contrary that there exists $j_0 \in [0, s-2]$ such that

$$\#\left(\begin{array}{c|c} j_0 & j_0+1 \\ \hline j_0 & j_0+1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} > 0. \tag{5.3}$$

We are going to show that

$$\#\left(\begin{array}{c|c} j_0 & j_0+1 \\ \hline j_0 & j_0+1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} > 0 \Longrightarrow \#\left(\begin{array}{c|c} j_0-1 & j_0 \\ \hline j_0-1 & j_0 \end{array}\right)_{\mathbb{P}_1/\mathbb{P}_{k+s-1}} > 0 \Longrightarrow \dots \Longrightarrow \#\left(\begin{array}{c|c} 0 & 1 \\ \hline 0 & 1 \end{array}\right)_{\mathbb{P}_j/\mathbb{P}_{k+s-1}} > 0,$$

which obviously contradicts

$$\#\left(\begin{array}{c|c} 0 & 1 \\ \hline 0 & 1 \end{array}\right)_{\mathbb{P}_j/\mathbb{P}_{k+s-1}} = 0.$$

By (5.3) there exists $t_0 \in \mathbb{P}/\mathbb{P}_{k+s-1}$ such that

$$r(D_{(s-1)\times(k-1)}(t_0)) = r(D_{s\times(k-1)}(t_0)) = j_0$$

and

$$r(D_{(s-1)\times k}(t_0)) = r(D_{s\times k}(t_0)) = j_0+1.$$

Consider the following partition of the matrix $D_{s\times k}(t_0)$

$$\left(\begin{array}{c|ccccc|c}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & & \alpha_k \\ \hline
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & & \vdots \\
\vdots & \vdots & \vdots & \vdots & \vdots & & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{k+s-3} & & \alpha_{k+s-2} \\ \hline
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{k+s-2} & & \alpha_{k+s-1}
\end{array}\right).$$

By (5.3), $r(D^2_{(s-1)\times(k-1)}(t_0)) \leq j_0$ and $r(D_{(s-1)\times k}(t_0)) = j_0 + 1$, it follows that $r(D^2_{(s-1)\times(k-1)}(t_0)) = j_0$.

Let $a_1, a_2, \ldots, a_k$ denote the columns of $D_{(s-1)\times k}(t_0)$, that is

$$a_i = \begin{pmatrix} \alpha_i \\ \alpha_{i+1} \\ \vdots \\ \alpha_{i+s-2} \end{pmatrix}$$

Since $r(D^2_{(s-1)\times(k-1)}(t_0)) = j_0$ *and* $r(D_{(s-1)\times k}(t_0)) = j_0 + 1$, we have $a_1 \notin$ *span* $\{a_2, a_3, \ldots, a_k\}$, therefore $r(D^2_{(s-1)\times(k-2)}(t_0)) = j_0 - 1$.

Now $r(D^2_{(s-1)\times(k-2)}(t_0)) = j_0 - 1, \quad r(D_{(s-1)\times(k-1)}(t_0)) = j_0 \quad$ *and* $\quad r(D_{s\times(k-1)}(t_0)) = j_0$.
From (5.2) we get $r(D^2_{s\times(k-2)}(t_0)) = j_0 - 1$. Thus we obtain

$r(D^2_{(s-1)\times(k-2)}(t_0)) = r(D^2_{s\times(k-2)}(t_0)) = j_0 - 1 \quad$ *and* $\quad r(D^2_{(s-1)\times(k-1)}(t_0)) = r(D^2_{s\times(k-1)}(t_0)) = j_0$.

Consider now the matrix

$$D^2_{s\times(k-1)}(t_0) = \left(\begin{array}{ccccc|c}
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_{k-1} & \alpha_k \\
\alpha_3 & \alpha_4 & \alpha_5 & \ldots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \ldots & \vdots & \vdots \\
\vdots & \vdots & \vdots & \ldots & \vdots & \vdots \\
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{k+s-3} & \alpha_{k+s-2} \\ \hline
\alpha_{s+1} & \alpha_{s+2} & \alpha_{s+2} & \ldots & \alpha_{k+s-2} & \alpha_{k+s-1}
\end{array}\right).$$

We get

$$\#\left(\begin{array}{c|c} j_0 - 1 & j_0 \\ \hline j_0 - 1 & j_0 \end{array}\right)_{\mathbb{P}_1/\mathbb{P}_{k+s-1}} > 0.$$

We now have proved that

$$\#\left(\begin{array}{c|c} j_0 & j_0+1 \\ \hline j_0 & j_0+1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} > 0 \Longrightarrow \#\left(\begin{array}{c|c} j_0-1 & j_0 \\ \hline j_0-1 & j_0 \end{array}\right)_{\mathbb{P}_1/\mathbb{P}_{k+s-1}} > 0.$$

We repeat this procedure and obtain after finitely many steps

$$\#\left(\begin{array}{c|c} 0 & 1 \\ \hline 0 & 1 \end{array}\right)_{\mathbb{P}_{j_0}/\mathbb{P}_{k+s-1}} > 0.$$

From

$$D^{j_0+1}_{s\times(k-j_0)}(t_0) = \left(\begin{array}{ccccc|c} \alpha_{j_0+1} & \alpha_{j_0+2} & \alpha_{j_0+3} & \ldots & \alpha_{k-1} & \alpha_k \\ \alpha_{j_0+2} & \alpha_{j_0+3} & \alpha_{j_0+4} & \ldots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \ldots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \ldots & \vdots & \vdots \\ \alpha_{s+j_0-1} & \alpha_{s+j_0} & \alpha_{s+j_0+1} & \ldots & \alpha_{k+s-3} & \alpha_{k+s-2} \\ \hline \\ \alpha_{s+j_0} & \alpha_{s+j_0+1} & \alpha_{s+j_0+1} & \ldots & \alpha_{k+s-2} & \alpha_{k+s-1} \end{array}\right),$$

we obviously get

$$\#\left(\begin{array}{c|c} 0 & 1 \\ \hline 0 & 1 \end{array}\right)_{\mathbb{P}_{j_0}/\mathbb{P}_{k+s-1}} = 0.$$

Arguing in the same way with the transpose of $D_{s\times k}(t)$, observing that the rank of a matrix is equal to the rank of its transpose, yields $\#\left(\begin{array}{c|c} j & j \\ \hline j+1 & j+1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} = 0$.

$\square$

Lemma 5.5. *We have*

$$\Gamma_i^{s\times(k-1)} = \Gamma_i^{(s-1)\times k} \quad \textit{if} \quad 0 \leq i \leq s-2, \quad s \leq k.$$

Proof. Consider the Matrix $D_{s\times k}(t)$

$$\left(\begin{array}{ccccc|c} \alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{k+s-3} & \alpha_{k+s-2} \\ \hline \\ \alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{k+s-2} & \alpha_{k+s-1} \end{array}\right).$$

By Lemmas 5.2, 5.3 and 5.4 we have

$$\begin{aligned}
2\cdot\Gamma_i^{s\times(k-1)} = {} & \#\left(\begin{array}{c|c} i & i \\ \hline i & i \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} + \#\left(\begin{array}{c|c} i & i \\ \hline i & i+1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} + \#\left(\begin{array}{c|c} i & i+1 \\ \hline i & i+1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} + \#\left(\begin{array}{c|c} i-1 & i-1 \\ \hline i & i-1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} \\
+ {} & \#\left(\begin{array}{c|c} i-1 & i-1 \\ \hline i & i \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} + \#\left(\begin{array}{c|c} i-1 & i \\ \hline i & i \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} + \#\left(\begin{array}{c|c} i-1 & i \\ \hline i & i+1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} \\
= {} & 2\cdot\#\left(\begin{array}{c|c} i & i \\ \hline i & i \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} + \#\left(\begin{array}{c|c} i-1 & i \\ \hline i & i+1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}
\end{aligned}$$

and

$$\begin{aligned}
2\cdot\Gamma_i^{(s-1)\times k} = {} & \#\left(\begin{array}{c|c} i & i \\ \hline i & i \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} + \#\left(\begin{array}{c|c} i & i \\ \hline i & i+1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} + \#\left(\begin{array}{c|c} i & i \\ \hline i+1 & i+1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} \\
+ {} & \#\left(\begin{array}{c|c} i-1 & i \\ \hline i-1 & i \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} + \#\left(\begin{array}{c|c} i-1 & i \\ \hline i & i \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} + \#\left(\begin{array}{c|c} i-1 & i \\ \hline i & i+1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} \\
= {} & 2\cdot\#\left(\begin{array}{c|c} i & i \\ \hline i & i \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} + \#\left(\begin{array}{c|c} i-1 & i \\ \hline i & i+1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}.
\end{aligned}$$

$\square$

Lemma 5.6. *We have (see Definition 1.1)*

$$\int_{\mathbb{P}} g^{2q}(t)dt = 2^{(s+k-2)(2q-1)}\cdot\sum_{j=0}^{s-1}\#\left(\begin{array}{c|c} j & j \\ \hline j & j \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}\cdot 2^{-2qj}.$$

Proof. From Lemma 5.2, by observing that g(t) is constants on cosets of $\mathbb{P}_{k+s-1}$, we obtain

$$\begin{aligned}
\int_{\mathbb{P}} g^{2q}(t)dt = {} & \sum_{j=0}^{s-1} 2^{(s+k-2-j)2q}\cdot\#\left(\begin{array}{c|c} j & j \\ \hline j & j \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}\int_{\mathbb{P}_{k+s-1}} dt \\
& + \sum_{j=0}^{s-1} 2^{(s+k-2-j)2q}\cdot\#\left(\begin{array}{c|c} j & j \\ \hline j & j+1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}\int_{\mathbb{P}_{k+s-1}} dt = \\
& 2\cdot\sum_{j=0}^{s-1} 2^{(s+k-2-j)2q}\cdot\#\left(\begin{array}{c|c} j & j \\ \hline j & j \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} 2^{-(k+s-1)} = \\
& 2^{(s+k-2)(2q-1)}\cdot\sum_{j=0}^{s-1}\#\left(\begin{array}{c|c} j & j \\ \hline j & j \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}\cdot 2^{-2qj}
\end{aligned}$$

which prove Lemma 5.6. $\square$

6. Exponential sums in $\mathbb{K}\times\mathbb{K}$

[See Definition 1.2]. In this section we compute exponential sums in $\mathbb{P}^2$, associated to a matrix of the form $\left[\frac{A}{b_-}\right]$, where A is a $(m+1)\times k$ persymmetic matrix over $\mathbb{F}_2$ and b_- a $1\times k$ matrix with entries in $\mathbb{F}_2$.

Lemma 6.1. *Set* $\eta = \sum_{j\geq 1}\beta_j T^{-j}\in\mathbb{P}$ *and* $Y = \sum_{i=1}^{k-1}\delta_i T^i\in\mathbb{F}_2[T]$, $\quad degY\leq k-1$.

Then

$$E(\eta Y) = \begin{cases} 1 & \textit{if } \sum_{j=1}^{k-1}\beta_j\delta_j = 0, \\ -1 & \textit{if } \sum_{j=1}^{k-1}\beta_j\delta_j = 1. \end{cases}$$

Proof. By lemmas 4.4, 4.5 with $t \longrightarrow \eta$ and s $= 1$ we have

$$\nu\left(\{\eta Y\}\right) > 1 \Longleftrightarrow Y \in \ker D_{1\times k}(\eta) \Longleftrightarrow \sum_{j=1}^{k-1}\beta_j\delta_j = 0,$$

$$\nu\left(\{\eta Y\}\right) = 1 \Longleftrightarrow \sum_{j=1}^{k-1}\beta_j\delta_j = 1.$$

□

Lemma 6.2. *Let* $(t,\eta) \in \mathbb{P}\times\mathbb{P}$ *and set*

$$g_{m,k}(t,\eta) = g(t,\eta) = \sum_{degY\leq k-1}\sum_{degZ\leq m} E(tYZ)\sum_{degU=0} E(\eta YU).$$

Then

$$g(t,\eta) = \begin{cases} 2^{k+m+1-r(D_{(1+m)\times k}(t))} & \textit{if } r(D_{(1+m)\times k}(t)) = r(D^{\left[\substack{1\\1+m}\right]\times k}(t,\eta)), \\ 0 & \textit{otherwise.} \end{cases}$$

Proof. Consider the matrix [see Definition 1.2]

$$D^{\left[\substack{1\\1+m}\right]\times k}(t,\eta) = \left(\begin{array}{cccccc} \alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{1+m} & \alpha_{2+m} & \alpha_{3+m} & \ldots & \alpha_{k+m-1} & \alpha_{k+m} \\ \hline \beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k \end{array}\right).$$

By Lemmas 4.4, 4.5 and 6.1 we obtain

$$\begin{aligned}
g(t,\eta) &= 2^{1+m} \sum_{\substack{degY\leq k-1\\ Y\in kerD_{(1+m)\times k}(t)}} E(\eta Y)\\
&= 2^{1+m}\cdot\Big[\sum_{\substack{degY\leq k-1\\ Y\in kerD^{\left[\begin{smallmatrix}1\\1+m\end{smallmatrix}\right]\times k}(t,\eta)}} 1 - \sum_{\substack{degY\leq k-1\\ Y\in kerD_{(1+m)\times k}(t)\\ \sum_{j=0}^{k-1}\beta_j\delta_j=1}} 1\Big]\\
&= 2^{1+m}\cdot\Big[\sum_{\substack{degY\leq k-1\\ Y\in kerD^{\left[\begin{smallmatrix}1\\1+m\end{smallmatrix}\right]\times k}(t,\eta)}} 1 - \Big(\sum_{\substack{degY\leq k-1\\ Y\in kerD_{(1+m)\times k}(t)}} 1 - \sum_{\substack{degY\leq k-1\\ Y\in kerD^{\left[\begin{smallmatrix}1\\1+m\end{smallmatrix}\right]\times k}(t,\eta)}} 1\Big)\Big]\\
&= 2^{1+m}\cdot\Big[2\cdot\sum_{\substack{degY\leq k-1\\ Y\in kerD^{\left[\begin{smallmatrix}1\\1+m\end{smallmatrix}\right]\times k}(t,\eta)}} 1 - \sum_{\substack{degY\leq k-1\\ Y\in kerD_{(1+m)\times k}(t)}} 1\Big]\\
&= 2^{1+m}\cdot\Big[2\cdot 2^{k-r(D^{\left[\begin{smallmatrix}1\\1+m\end{smallmatrix}\right]\times k}(t,\eta))} - 2^{k-r(D_{(1+m)\times k}(t))}\Big].
\end{aligned}$$

□

Lemma 6.3. *Set* $h(t)=\sum_{deg\leq k-1}\sum_{degZ\leq m}E(tYZ)=h_{m+1,k}(t)$ *and let* $q\geq 2$ *be an integer, then*

$$g^q(t,\eta)=g(t,\eta)\cdot h^{q-1}(t).$$

Proof. Obviously we get

$$g^2(t,\eta)=\Big[2^{1+m}\sum_{\substack{degY_1\leq k-1\\ Y_1\in kerD_{(1+m)\times k}(t)}}E(\eta Y_1)\Big]\cdot\Big[2^{1+m}\sum_{\substack{degY_2\leq k-1\\ Y_2\in kerD_{(1+m)\times k}(t)}}E(\eta Y_2)\Big].$$

Define

$$\begin{cases} Y_1+Y_2=Y_3 & degY_3\leq k-1\\ Y_1=Y_4 & degY_4\leq k-1.\end{cases}$$

Then we obtain

$$\begin{aligned}
g^2(t,\eta) &= 2^{2(1+m)}\sum_{\substack{degY_1\leq k-1\\ Y_1\in kerD_{(1+m)\times k}(t)}}\ \sum_{\substack{degY_2\leq k-1\\ Y_2\in kerD_{(1+m)\times k}(t)}}E(\eta(Y_1+Y_2))\\
&= \Big[2^{1+m}\sum_{\substack{degY_4\leq k-1\\ Y_4\in kerD_{(1+m)\times k}(t)}}1\Big]\cdot\Big[2^{1+m}\sum_{\substack{degY_3\leq k-1\\ Y_3\in kerD_{(1+m)\times k}(t)}}E(\eta Y_3)\Big]\\
&= h(t)\cdot g(t,\eta).
\end{aligned}$$

By recurrence on q we get $g^q(t,\eta)=g(t,\eta)\cdot h^{q-1}(t)$. □

Lemma 6.4. *We have*

$$\int_{\mathbb{P}}\int_{\mathbb{P}}g^q(t,\eta)dtd\eta = 2^{q(k+m+1)-2k-m}\sum_{i=0}^{inf(k,1+m)}\sigma_{i,i}^{\left[\begin{smallmatrix}1\\1+m\end{smallmatrix}\right]\times k}2^{-iq}.$$

Proof. Recall Definition 1.2, then Lemma 6.2 gives, by observing that $g(t,\eta)$ is constant on cosets of
$\mathbb{P}_{k+m} \times \mathbb{P}_k$.

$$\begin{aligned}\int_{\mathbb{P}}\int_{\mathbb{P}} g^q(t,\eta)dtd\eta &= \sum_{\substack{(t,\eta)\in\mathbb{P}/\mathbb{P}_{k+m}\times\mathbb{P}/\mathbb{P}_k \\ r(D_{(1+m)\times k}(t))=r(D^{\left[\genfrac{}{}{0pt}{}{1}{1+m}\right]\times k}(t,\eta))}} 2^{q(k+m+1-r(D_{(1+m)\times k}(t))}\int_{\mathbb{P}_{k+m}} dt \int_{\mathbb{P}_k} d\eta \\ &= \sum_{i=0}^{inf(k,1+m)} \sigma_{i,i}^{\left[\genfrac{}{}{0pt}{}{1}{1+m}\right]\times k} 2^{q(k+m+1-i)}2^{-(k+m)}2^{-k}.\end{aligned}$$

□

Lemma 6.5. *Equally*

$$\int_{\mathbb{P}}\int_{\mathbb{P}} g^q(t,\eta)dtd\eta = 2^{q(k+m+1)-2k-m}\sum_{i=0}^{inf(k,1+m)} 2^i\Gamma_i^{(1+m)\times k}2^{-iq}.$$

Proof. From Lemma 6.3 and Lemma 4.6 with s = m+1 we get by Fubini's theorem

$$\begin{aligned}\int_{\mathbb{P}}\int_{\mathbb{P}} g^q(t,\eta)dtd\eta &= \int_{\mathbb{P}}\int_{\mathbb{P}} g(t,\eta)h^{q-1}(t)dtd\eta = \int_{\mathbb{P}} h^{q-1}(t)\big(\int_{\mathbb{P}} g(t,\eta)d\eta\big)dt = 2^{1+m}\int_{\mathbb{P}} h^{q-1}(t)dt \\ &= 2^{1+m}\cdot \sum_{t\in\mathbb{P}/\mathbb{P}_{k+m}} 2^{[k+m+1-r(D_{(m+1)\times k}(t))]\cdot(q-1)}\int_{\mathbb{P}_{k+m}} dt \\ &= 2^{1+m}\cdot \sum_{i=0}^{inf(k,1+m)} \Gamma_i^{(m+1)\times k}\cdot 2^{(m+1+k-i)\cdot(q-1)}\int_{\mathbb{P}_{k+m}} dt \\ &= 2^{1+m}\cdot 2^{(q-2)(k+m+1)+1}\sum_{i=0}^{inf(k,1+m)} \Gamma_i^{(1+m)\times k}2^{-(q-1)i}.\end{aligned}$$

□

Lemma 6.6. *For all* $0 \leq i \leq inf(k,1+m)$ *we have* $\sigma_{i,i}^{\left[\genfrac{}{}{0pt}{}{1}{1+m}\right]\times k} = 2^i\Gamma_i^{(1+m)\times k}$.

Proof. Lemma 6.4 and Lemma 6.5 give

$$\sum_{i=0}^{inf(k,1+m)} \big(\sigma_{i,i}^{\left[\genfrac{}{}{0pt}{}{1}{1+m}\right]\times k} - 2^i\Gamma_i^{(1+m)\times k}\big)\cdot 2^{-iq} = 0 \quad for\ all\ q \geq 2.$$

□

Lemma 6.7. *For all* $1 \leq i \leq inf(k,2+m)$ *we have*
$\sigma_{i-1,i}^{\left[\genfrac{}{}{0pt}{}{1}{1+m}\right]\times k} = (2^k - 2^{i-1})\Gamma_{i-1}^{(1+m)\times k}$.

Proof. Consider the matrix

$$D^{\left[\substack{1\\1+m}\right]\times k}(t,\eta) = \left(\begin{array}{cccccc} \alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{1+m} & \alpha_{2+m} & \alpha_{3+m} & \ldots & \alpha_{k+m-1} & \alpha_{k+m} \\ \hline \beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k \end{array}\right).$$

Clearly Lemma 6.6 yields

$$\begin{aligned} 2^k \cdot \Gamma_{i-1}^{(1+m)\times k} &= \sigma_{i-1,i-1}^{\left[\substack{1\\1+m}\right]\times k} + \sigma_{i-1,i}^{\left[\substack{1\\1+m}\right]\times k} \\ \Leftrightarrow \sigma_{i-1,i}^{\left[\substack{1\\1+m}\right]\times k} &= 2^k \cdot \Gamma_{i-1}^{(1+m)\times k} - 2^{i-1} \cdot \Gamma_{i-1}^{(1+m)\times k} \\ \Leftrightarrow \sigma_{i-1,i}^{\left[\substack{1\\1+m}\right]\times k} &= (2^k - 2^{i-1}) \cdot \Gamma_{i-1}^{(1+m)\times k} \qquad \text{if } 1 \leq i \leq inf(k, 2+m) \end{aligned}$$

□

Lemma 6.8. *For all* $0 \leq i \leq inf(k, 2+m)$ *we have*
$\Gamma_i^{\left[\substack{1\\1+m}\right]\times k} = (2^k - 2^{i-1}) \cdot \Gamma_{i-1}^{(1+m)\times k} + 2^i \Gamma_i^{(1+m)\times k}$.

Proof. From Lemmas 6.6, 6.7 we get

$$\Gamma_i^{\left[\substack{1\\1+m}\right]\times k} = \sigma_{i,i}^{\left[\substack{1\\1+m}\right]\times k} + \sigma_{i-1,i}^{\left[\substack{1\\1+m}\right]\times k} = 2^i \Gamma_i^{(1+m)\times k} + (2^k - 2^{i-1}) \cdot \Gamma_{i-1}^{(1+m)\times k}.$$

□

Lemma 6.9. *Set* $f(t,\eta) = \sum_{degY \leq k-1} \sum_{degZ \leq m} E(tYZ) \sum_{degU \leq 0} E(\eta YU)$
for $k \geq 1$ *and* $m \geq 0$.
Then

$$f(t,\eta) = 2^{k+m+2-r(r(D^{\left[\substack{1\\1+m}\right]\times k}(t,\eta))}$$

and

$$\int_{\mathbb{P}} \int_{\mathbb{P}} f^q(t,\eta) dt d\eta = 2^{q(k+m+2)-2k-m} \sum_{i=0}^{inf(k,2+m)} \Gamma_i^{\left[\substack{1\\1+m}\right]\times k} 2^{-iq}.$$

Proof. By using Lemma 4.4 and Lemma 4.5 we obtain

$$\begin{aligned}
f(t,\eta) &= \sum_{degY\leq k-1}\sum_{degZ\leq m} E(tYZ)\sum_{degU\leq 0} E(\eta YU) \\
&= 2^{2+m}\sum_{\substack{degY\leq k-1 \\ Y\in kerD_{(1+m)\times k}(t) \\ Y\in kerD_{1\times k}(\eta)}} 1 \\
&= 2^{2+m}\sum_{\substack{degY\leq k-1 \\ Y\in kerD^{\left[\substack{1\\1+m}\right]\times k}(t,\eta)}} 1 = 2^{k+m+2-r(r(D^{\left[\substack{1\\1+m}\right]\times k}(t,\eta)}.
\end{aligned}$$

Then we have, by observing that $f(t,\eta)$ is constant on cosets of $\mathbb{P}_{k+m}\times\mathbb{P}_k$

$$\begin{aligned}
\int_{\mathbb{P}}\int_{\mathbb{P}} f^q(t,\eta)dtd\eta &= \sum_{(t,\eta)\in\mathbb{P}/\mathbb{P}_{k+m}\times\mathbb{P}/\mathbb{P}_k} 2^{q(k+m+2-r(r(D^{\left[\substack{1\\1+m}\right]\times k}(t,\eta))}\int_{\mathbb{P}_{k+m}} dt\int_{\mathbb{P}_k} d\eta \\
&= \sum_{i=0}^{inf(k,2+m)} \Gamma_i^{\left[\substack{1\\1+m}\right]\times k} 2^{q(k+m+2-i)}2^{-(k+m)}2^{-k}.
\end{aligned}$$

□

7. Proof of Theorem 3.1

We prove Theorem 3.1 by induction on s (recall that $s\leq k$).
Set $h_{s,k}(t) = \sum_{degY\leq k-1}\sum_{degZ\leq s-1} E(tYZ)$. Let s = 2 and consider the matrix $D_{2\times k}(t)$, that is

$$D_{2\times k}(t) = \begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \ldots\alpha_k \\ \alpha_2 & \alpha_3 & \alpha_4 & \ldots\alpha_{k+1} \end{pmatrix}.$$

We have

$$\sum_{i=0}^{2}\Gamma_i^{2\times k} = 2^{k+1}. \tag{7.1}$$

By Lemma 4.6 with s = 2 and q = 1

$$\int_{\mathbb{P}} h_{2,k}(t)dt = \int_{\mathbb{P}} 2^{k+2-r(D_{2\times k}(t))}dt = 2\cdot\sum_{i=0}^{2}\Gamma_i^{2\times k}\cdot 2^{-i}. \tag{7.2}$$

On the other hand

(7.3)

$$\int_{\mathbb{P}} h_{2,k}(t)dt = Card\left\{(Y,Z), degY\leq k-1, degZ\leq 1 \mid Y\cdot Z = 0\right\} = 2^k{+}2^2{-}1.$$

From (7.2) and (7.3) we obtain

$$2\Gamma_1^{2\times k} + \Gamma_2^{2\times k} = 2^{k+1} + 2. \tag{7.4}$$

From (7.1) and (7.4) we deduce

$$\Gamma_i^{2\times k} = \begin{cases} 1 & if \quad i=0, \\ 3 & if \quad i=1, \\ 2^{(k+1)}-4 & if \quad i=2. \end{cases}$$

Let s = 3 and consider the matrix $D_{3\times k}(t)$

$$D_{3\times k}(t) = \begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \ldots\alpha_k \\ \alpha_2 & \alpha_3 & \alpha_4 & \ldots\alpha_{k+1} \\ \alpha_3 & \alpha_4 & \alpha_5 & \ldots\alpha_{k+2} \end{pmatrix}.$$

We have

$$\sum_{i=0}^{3}\Gamma_i^{3\times k} = 2^{k+2}. \tag{7.5}$$

By Lemma 5.5

$$\Gamma_i^{3\times k} = \Gamma_i^{2\times(k+1)} \; if \; 0\leq i\leq 1, \quad k\geq 3. \tag{7.6}$$

Using (7.5) and (7.6), we get

$$\Gamma_2^{3\times k} + \Gamma_3^{3\times k} = 2^{k+2}-2^2. \tag{7.7}$$

By Lemma 4.6 with s = 3 and q =1

$$\int_{\mathbb{P}} h_{3,k}(t)dt = \int_{\mathbb{P}} 2^{k+3-r(D_{3\times k}(t))}dt = 2\cdot\sum_{i=0}^{3}\Gamma_i^{3\times k}\cdot 2^{-i}. \tag{7.8}$$

On the other hand

(7.9)

$$\int_{\mathbb{P}} h_{3,k}(t)dt = Card\left\{(Y,Z), degY\leq k-1, degZ\leq 2 \mid Y\cdot Z=0\right\} = 2^k+2^3-1.$$

From (7.8) and (7.9) we obtain

$$2\Gamma_2^{3\times k} + \Gamma_3^{3\times k} = 2^{k+2}+2^3. \tag{7.10}$$

From (7.7) and (7.10) we deduce

$$\Gamma_i^{3\times k} = \begin{cases} 1 & if \quad i=0, \\ 3 & if \quad i=1, \\ 12 & if \quad i=2, \\ 2^{(k+2)}-2^4 & if \quad i=3. \end{cases}$$

Let s = 4. In the same way we obtain

$$\Gamma_i^{4\times k} = \begin{cases} 1 & if \quad i=0, \\ 3 & if \quad i=1, \\ 12 & if \quad i=2, \\ 48 & if \quad i=3, \\ 2^{(k+3)}-2^6 & if \quad i=4. \end{cases}$$

Assume that for $k \geq s-1$ we have

$$(H) \quad \Gamma_{i}^{(s-1)\times k} = \begin{cases} 1 & if \quad i = 0, \\ 3\cdot 2^{2(i-1)} & if \quad 1 \leq i \leq s-2, \\ 2^{(k+s-2)} - 2^{2s-4} & if \quad i = s-1. \end{cases}$$

From (H) and Lemma 5.5 it follows

$$\Gamma_{i}^{s\times(k-1)} = \Gamma_{i}^{(s-1)\times k} = 3\cdot 2^{2(i-1)} \; if \; 0 \leq i \leq s-2 \quad , k \geq s+1. \tag{7.11}$$

Consider the following partition of the matrix $D_{s\times k}(t)$

$$\left(\begin{array}{ccccc|c}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{k+s-3} & \alpha_{k+s-2} \\
\hline
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{k+s-2} & \alpha_{k+s-1}
\end{array}\right).$$

By (7.11) we get

$$\sum_{i=0}^{s-2} \Gamma_{i}^{s\times(k-1)} = \sum_{i=0}^{s-2} 3\cdot 2^{2(i-1)} = 2^{2s-4} \tag{7.12}$$

which implies

$$\Gamma_{s-1}^{s\times(k-1)} + \Gamma_{s}^{s\times(k-1)} = 2^{k+s-2} - 2^{2s-4}. \tag{7.13}$$

From Lemma 4.6 it follows with $k \longrightarrow k-1$ and q = 1

$$\int_{\mathbb{P}} h_{s,k-1}(t)dt = \int_{\mathbb{P}} 2^{k+s-1-r(D_{s\times(k-1)}(t))}dt = 2\cdot\sum_{i=0}^{s} \Gamma_{i}^{s\times(k-1)}\cdot 2^{-i}. \tag{7.14}$$

On the other hand

(7.15)

$$\int_{\mathbb{P}} h_{s,k-1}(t)dt = Card\left\{(Y,Z), degY \leq k-2, degZ \leq s-1 \mid Y\cdot Z = 0\right\} = 2^{k-1}+2^{s}-1.$$

From (**??**) and (7.15) we obtain

$$\sum_{i=0}^{s} \Gamma_{i}^{s\times(k-1)}\cdot 2^{-i} = 2^{k-2} + 2^{s-1} - 2^{-1}. \tag{7.16}$$

In view of (7.11)

$$\sum_{i=0}^{s-2} \Gamma_{i}^{s\times(k-1)}\cdot 2^{-i} = \sum_{i=0}^{s-2} 3\cdot 2^{2(i-1)}\cdot 2^{-i} = 3\cdot 2^{s-3} - 2^{-1}. \tag{7.17}$$

From (7.16) and (7.17)

$$2\Gamma_{s-1}^{s\times(k-1)} + \Gamma_{s}^{s\times(k-1)} = 2^{k+s-2} + 2^{2s-3}. \tag{7.18}$$

By (7.11), (7.13) and (7.18) we deduce for $k \geq s+1$

$$\Gamma_i^{s\times(k-1)} = \begin{cases} 1 & if \quad i=0, \\ 3\cdot 2^{2(i-1)} & if \quad 1 \leq i \leq s-1, \\ 2^{(k+s-2)} - 2^{2s-2} & if \quad i = s. \end{cases}$$

□

8. Proof of Theorem 3.3

The proof of Theorem 3.3 is based on the relation between Theorem 3.1 and the following Lemmas.

Lemma 8.1. *Let $t \in \mathbb{P}$, $(s \leq k)$ then*

(8.1)

$$\Gamma_i^{s\times k} = \begin{cases} \#\left(\begin{array}{c|c} 0 & 0 \\ \hline 0 & 0 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} & \textit{if } i=0, \\ \#\left(\begin{array}{c|c} 1 & 1 \\ \hline 1 & 1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} + \#\left(\begin{array}{c|c} 0 & 0 \\ \hline 0 & 0 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} & \textit{if } i=1, \\ \#\left(\begin{array}{c|c} i & i \\ \hline i & i \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} + \#\left(\begin{array}{c|c} i-1 & i-1 \\ \hline i-1 & i-1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} + \#\left(\begin{array}{c|c} i-2 & i-1 \\ \hline i-1 & i \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} & \textit{if } 2 \leq i \leq s-1, \\ \#\left(\begin{array}{c|c} s-1 & s-1 \\ \hline s-1 & s-1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} + \#\left(\begin{array}{c|c} s-1 & s-1 \\ \hline s & s \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} + \#\left(\begin{array}{c|c} s-2 & s-1 \\ \hline s-1 & s \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} & \textit{if } i=s \end{cases}$$

and

(8.2)

$$2\cdot\Gamma_{i-1}^{s\times(k-1)} = \begin{cases} 2\cdot\#\left(\begin{array}{c|c} 0 & 0 \\ \hline 0 & 0 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} & \textit{if } i=1, \\ 2\cdot\#\left(\begin{array}{c|c} i-1 & i-1 \\ \hline i-1 & i-1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} + \#\left(\begin{array}{c|c} i-2 & i-1 \\ \hline i-1 & i \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} & \textit{if } 2 \leq i \leq s-1, \\ 2\cdot\#\left(\begin{array}{c|c} s-1 & s-1 \\ \hline s-1 & s-1 \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} + \#\left(\begin{array}{c|c} s-2 & s-1 \\ \hline s-1 & s \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} & \textit{if } i=s, \\ \#\left(\begin{array}{c|c} s-1 & s-1 \\ \hline s & s \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} & \textit{if } i=s+1. \end{cases}$$

Proof. It suffices to reproduce carefully the proof of Lemma 5.5. □

Lemma 8.2. *Let $t \in \mathbb{P}$, $(s \leq k)$ then*

(8.3)
$$\Gamma_{i}^{s\times k}-2\cdot\Gamma_{i-1}^{s\times(k-1)}=\begin{cases}\#\left(\begin{array}{c|c}1&1\\\hline 1&1\end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}-\#\left(\begin{array}{c|c}0&0\\\hline 0&0\end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} & \textit{if } i=1,\\ \#\left(\begin{array}{c|c}i&i\\\hline i&i\end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}-\#\left(\begin{array}{c|c}i-1&i-1\\\hline i-1&i-1\end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} & \textit{if } 2\le i\le s-1,\\ \#\left(\begin{array}{c|c}s-1&s-1\\\hline s&s\end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}-\#\left(\begin{array}{c|c}s-1&s-1\\\hline s-1&s-1\end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} & \textit{if } i=s.\end{cases}$$

Proof. Follows from (8.1) and (8.2). ☐

Lemma 8.3. *We have*

$$\Gamma_{i}^{s\times k}-2\cdot\Gamma_{i-1}^{s\times(k-1)}=\begin{cases}1 & \textit{if } i=1,\\ 3\cdot 2^{2i-3} & \textit{if } 2\le i\le s-1,\\ 2^{k+s-1}-5\cdot 2^{2s-3} & \textit{if } i=s.\end{cases}\tag{8.4}$$

Proof. The assertion follows immediately from Theorem 3.1. ☐

Theorem 3.1 and Lemmas 8.1, 8.2, 8.3 give the following relations

$$\#\left(\begin{array}{c|c}1&1\\\hline 1&1\end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}-\#\left(\begin{array}{c|c}0&0\\\hline 0&0\end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}=1,\tag{8.5}$$

$$\#\left(\begin{array}{c|c}i&i\\\hline i&i\end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}-\#\left(\begin{array}{c|c}i-1&i-1\\\hline i-1&i-1\end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}=3\cdot 2^{2i-3}\quad\text{if}\quad 2\le i\le s-1\tag{8.6}$$

(8.7)
$$\#\left(\begin{array}{c|c}s-1&s-1\\\hline s&s\end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}-\#\left(\begin{array}{c|c}s-1&s-1\\\hline s-1&s-1\end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}=2^{k+s-1}-5\cdot 2^{2s-3},$$

(8.8)
$$\#\left(\begin{array}{c|c}i&i\\\hline i&i\end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}+\#\left(\begin{array}{c|c}i-1&i-1\\\hline i-1&i-1\end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}+\#\left(\begin{array}{c|c}i-2&i-1\\\hline i-1&i\end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}=3\cdot 2^{2i-2}\quad\text{if}\quad 1\le i\le s-1,$$

(8.9)
$$\#\left(\begin{array}{c|c}s-1&s-1\\\hline s-1&s-1\end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}+\#\left(\begin{array}{c|c}s-1&s-1\\\hline s&s\end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}+\#\left(\begin{array}{c|c}s-2&s-1\\\hline s-1&s\end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}=2^{k+s-1}-2^{2s-2}.$$

Now we proceed as follows:

First we deduce by (8.5) and (8.6)

$$\#\left(\begin{array}{c|c}i&i\\\hline i&i\end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}=2^{2i-1}\quad \textit{if } 1\le i\le s-1.\tag{8.10}$$

Secondly by (8.8) and (8.10)

$$\#\left(\begin{array}{c|c}i-2&i-1\\\hline i-1&i\end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}}=2^{2i-3}\quad \textit{if } 2\le i\le s-1.$$

Thirdly by (8.7) and (8.10) with i = s - 1

$$\#\left(\begin{array}{c|c} s-1 & s-1 \\ \hline s & s \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} = 2^{k+s-1} - 2^{2s-1}. \tag{8.11}$$

Last by (8.9), (8.11) and (8.10) with i = s - 1

$$\#\left(\begin{array}{c|c} s-2 & s-1 \\ \hline s-1 & s \end{array}\right)_{\mathbb{P}/\mathbb{P}_{k+s-1}} = 2^{2s-3}.$$

9. Proofs of Theorems 3.4,3.5,3.6,3.7 and 3.8

9.1. **Proof of Theorem 3.4.** Apply Lemma 4.6 and note that

$$\int_{\mathbb{P}} h^q(t)dt = \int_{\mathbb{P}} \Big[\sum_{degY\leq k-1}\ \sum_{degZ\leq s-1} E(tYZ)\Big]^q dt =$$

$$\int_{\mathbb{P}} \sum_{Y_1}\sum_{Z_1}\sum_{Y_2}\sum_{Z_2}\cdots\sum_{Y_q}\sum_{Z_q} E(tY_1Z_1)\cdots E(tY_qZ_q)dt =$$

$$\sum_{Y_1}\sum_{Z_1}\sum_{Y_2}\sum_{Z_2}\cdots\sum_{Y_q}\sum_{Z_q}\int_{\mathbb{P}} E(t(Y_1Z_1+\cdots+Y_qZ_q))dt = R$$

$\square$

9.2. **Proof of Theorem 3.5.** Follows from Lemma 4.10 and Lemma 5.6. $\square$

9.3. **Proof of Theorem 3.6.** Follows from Lemma 6.2 and Lemma 6.4. $\square$

9.4. **Proof of Theorem 3.7.** Immediately from Lemma 6.9 . $\square$

9.5. **Proof of Theorem 3.8.** Let $2 \leq m \leq k-2$. Then, by Lemma 6.8 and Theorem 3.1 with s = m+1

$$
\begin{aligned}
\Gamma_0^{\left[\substack{1\\1+m}\right]\times k} &= 1,\\
\Gamma_1^{\left[\substack{1\\1+m}\right]\times k} &= \sigma_{0,1}+\sigma_{1,1} = (2^k-1)+2\cdot 3 = 2^k+5,\\
\Gamma_i^{\left[\substack{1\\1+m}\right]\times k} &= \sigma_{i-1,i}+\sigma_{i,i} = (2^k-2^{i-1})\cdot\Gamma_{i-1}^{(1+m)\times k}+2^i\cdot\Gamma_i^{(1+m)\times k}\\
&= (2^k-2^{i-1})\cdot 3\cdot 2^{2(i-2)}+2^i\cdot 3\cdot 2^{2(i-1)}) = 3\cdot 2^{k+2i-4}+21\cdot 2^{3i-5} \qquad \text{if } 2\leq i\leq m,\\
\Gamma_{m+1}^{\left[\substack{1\\1+m}\right]\times k} &= \sigma_{m,m+1}+\sigma_{m+1,m+1} = (2^k-2^m)\cdot\Gamma_{i-1}^{m\times k}+2^{m+1}\cdot\Gamma_{m+1}^{(1+m)\times k}\\
&= (2^k-2^m)\cdot 3\cdot 2^{2(m-1)}+2^{1+m}\cdot(2^{k+m}-2^{2m}) = 11\cdot[2^{k+2m-2}-2^{3m-2}],\\
\Gamma_{m+2}^{\left[\substack{1\\1+m}\right]\times k} &= \sigma_{m+1,m+2} = (2^k-2^{1+m})\cdot\Gamma_{m+1}^{(1+m)\times k} = (2^k-2^{1+m})\cdot(2^{k+m}-2^{2m})\\
&= 2^{2k+m}-3\cdot 2^{k+2m}+2^{3m+1}.
\end{aligned}
$$

The proof in the case $3\leq k\leq 1+m$ is similar. □

10. Proofs of Theorem 3.9 , Corollary 3.10 and Theorem 3.11

10.1. $\Gamma_i^{\left[\substack{n\\1+m}\right]\times k}$ written as a linear combination of the $\Gamma_{i-j}^{(1+m)\times k}$ for $0\leq j\leq n$..

Lemma 10.1. *[See Definitions 1.1, 1.2]. The number* $\Gamma_i^{\left[\substack{n\\1+m}\right]\times k}$, *expressed as a linear combination of the* $\Gamma_{i-j}^{(1+m)\times k}$ *for* $j=0,1,\ldots n$, *is given by the following recurrence formula*

(10.1)

$$
\Gamma_i^{\left[\substack{n\\1+m}\right]\times k} = \sum_{j=0}^{n}\Big[2^{(n-j)\cdot(i-j)}a_j^{(n)}\prod_{l=1}^{j}(2^k-2^{i-l})\Big]\cdot\Gamma_{i-j}^{(1+m)\times k}, n=1,2,\ldots \quad for \quad 0\leq i\leq inf(k,n+m+1),
$$

where $a_j^{(n)}$ *satisfies the linear recurrence relation*

$$
a_j^{(n)} = 2^j\cdot a_j^{(n-1)}+a_{j-1}^{(n-1)}, \quad n=2,3,4,\ldots \quad for \quad 1\leq j\leq n-1. \tag{10.2}
$$

We set

$$
\begin{aligned}
&a_0^{(n)} = a_n^{(n)} = 1\\
\text{and}\quad &\Gamma_{i-j}^{(1+m)\times k} = 0 \quad if \quad i-j\notin\{0,1,2,\ldots,inf(k,1+m)\}.
\end{aligned}
$$

We can then compute successively $a_j^{(n)}, \quad n \geq j \quad for \quad j = 1, 2, \ldots$ *and we obtain*

$$\begin{aligned} a_1^{(n)} &= 2^n - 1 \quad for \quad n \geq 1, \\ a_2^{(n)} &= \frac{2^{2n-1} - 3 \cdot 2^{n-1} + 1}{3} \quad for \quad n \geq 2, \\ a_j^{(n)} &= 2^{nj-j^2} + \sum_{l=0}^{n-j-1} a_{j-1}^{(n-1-l)} \cdot (2^j)^l \quad for \quad n \geq j. \end{aligned}$$

Proof. We prove Lemma 10.1 by induction on n.
A similar proof of Lemma 6.8 gives the following generalization
(10.3)

$$\Gamma_i^{\left[\begin{smallmatrix} n \\ 1+m \end{smallmatrix}\right]\times k} = 2^i \cdot \Gamma_i^{\left[\begin{smallmatrix} n-1 \\ 1+m \end{smallmatrix}\right]\times k} + (2^k - 2^{i-1}) \cdot \Gamma_{i-1}^{\left[\begin{smallmatrix} n-1 \\ 1+m \end{smallmatrix}\right]\times k} \quad for \quad 0 \leq i \leq inf(k, n+m+1).$$

From Lemma 6.8 and (10.3) with n=2 we get respectively for n =1 and n = 2

(10.4)

$$\Gamma_i^{\left[\begin{smallmatrix} 1 \\ 1+m \end{smallmatrix}\right]\times k} = 2^i \Gamma_i^{(1+m)\times k} + (2^k - 2^{i-1}) \cdot \Gamma_{i-1}^{(1+m)\times k} \quad for \quad 0 \leq i \leq inf(k, 2+m)$$

(10.5)

$$\Gamma_i^{\left[\begin{smallmatrix} 2 \\ 1+m \end{smallmatrix}\right]\times k} = 2^i \Gamma_i^{\left[\begin{smallmatrix} 1 \\ 1+m \end{smallmatrix}\right]\times k} + (2^k - 2^{i-1}) \cdot \Gamma_{i-1}^{\left[\begin{smallmatrix} 1 \\ 1+m \end{smallmatrix}\right]\times k} \quad for \quad 0 \leq i \leq inf(k, 3+m).$$

From (10.4) and (10.5) we deduce

(10.6)

$$\begin{aligned} \Gamma_i^{\left[\begin{smallmatrix} 2 \\ 1+m \end{smallmatrix}\right]\times k} &= 2^{2i} \Gamma_i^{(1+m)\times k} + 3 \cdot 2^{i-1}(2^k - 2^{i-1}) \cdot \Gamma_{i-1}^{(1+m)\times k} \\ &+ (2^k - 2^{i-1})(2^k - 2^{i-2}) \cdot \Gamma_{i-2}^{(1+m)\times k} \quad for \quad 0 \leq i \leq inf(k, 3+m). \end{aligned}$$

Hence by (10.4) and (10.6) the formula (10.1) holds for n = 1 and n = 2.
Assume now that (10.1) holds for the number n-1, that is

$$(10.7) \quad 2^i \cdot \Gamma_i^{\left[\begin{smallmatrix} n-1 \\ 1+m \end{smallmatrix}\right]\times k} = \sum_{j=0}^{n-1} 2^{(n-1-j)\cdot(i-j)} 2^i a_j^{(n-1)} \prod_{l=1}^{j} (2^k - 2^{i-l}) \cdot \Gamma_{i-j}^{(1+m)\times k},$$

(10.8)

$$(2^k-2^{i-1})\cdot\Gamma_{i-1}^{\left[\substack{n-1\\1+m}\right]\times k} = (2^k-2^{i-1})\big[\sum_{j=0}^{n-1}2^{(n-1-j)\cdot(i-1-j)}a_j^{(n-1)}\prod_{l=1}^{j}(2^k-2^{i-1-l})\cdot\Gamma_{i-1-j}^{(1+m)\times k}\big]$$

$$=\sum_{j=0}^{n-1}2^{(n-1-j)\cdot(i-1-j)}a_j^{(n-1)}\prod_{l=1}^{j+1}(2^k-2^{i-l})\cdot\Gamma_{i-(j+1)}^{(1+m)\times k}$$

$$=\sum_{j=1}^{n}2^{(n-j)\cdot(i-j)}a_{j-1}^{(n-1)}\prod_{l=1}^{j}(2^k-2^{i-l})\cdot\Gamma_{i-j}^{(1+m)\times k}.$$

From (10.7), (10.8) , and (10.2) it follows

(10.9)

$$\Gamma_i^{\left[\substack{n\\1+m}\right]\times k} = 2^{ni}\Gamma_i^{(1+m)\times k}+\sum_{j=1}^{n-1}2^{(n-j)\cdot(i-j)}(2^j a_j^{(n-1)}+a_{j-1}^{(n-1)})\prod_{l=1}^{j}(2^k-2^{i-l})\cdot\Gamma_{i-j}^{(1+m)\times k}$$

$$+\,a_{n-1}^{(n-1)}\prod_{l=1}^{n}(2^k-2^{i-l})\Gamma_{i-n}^{(1+m)\times k}$$

$$=\sum_{j=0}^{n}2^{(n-j)\cdot(i-j)}a_j^{(n)}\prod_{l=1}^{j}(2^k-2^{i-l})\cdot\Gamma_{i-j}^{(1+m)\times k}.$$

□

10.2. **Computation of a_i^n for $1 \leq i \leq n-1$.**

Lemma 10.2. *We have for* $1 \leq i \leq n-1$

$$a_i^{(n)} = \sum_{s=0}^{i-1}(-1)^s\prod_{l=0}^{i-(s+1)}\frac{2^{n+1}-2^l}{2^{i-s}-2^l}\cdot 2^{s(n-i)+\frac{s(s+1)}{2}}+(-1)^i\cdot 2^{in-\frac{i(i-1)}{2}}.$$

Proof. (See Definition 1.2) We recall that $\Gamma_i^{\left[\substack{n\\1}\right]\times k}$ denotes the number of $(n+1)\times k$ matrices over $\mathbb{F}_2$ of rank i.
By formula (10.1) with m = 0, we obtain

(10.10)

$$\Gamma_i^{\left[\substack{n\\1}\right]\times k} = \sum_{j=0}^{n}2^{(n-j)\cdot(i-j)}a_j^{(n)}\prod_{l=1}^{j}(2^k-2^{i-l})\cdot\Gamma_{i-j}^{1\times k}, n=1,2,\ldots \quad for \quad 0\leq i\leq inf(k,n+1).$$

We have obviously

$$\Gamma_{i-j}^{1\times k} = \begin{cases}1 & \text{if } i-j=0,\\ 2^k-1 & \text{if } i-j=1.\end{cases} \tag{10.11}$$

From (10.10) and (10.11) we deduce

(10.12)

$$\begin{aligned}
\Gamma_i^{\left[\begin{smallmatrix} n\\ 1\end{smallmatrix}\right]\times k} &= \sum_{j=i-1}^{i} 2^{(n-j)\cdot(i-j)} a_j^{(n)} \prod_{l=1}^{j}(2^k-2^{i-l})\cdot \Gamma_{i-j}^{1\times k} \\
&= a_{i-1}^{(n)}\cdot 2^{n-(i-1)}\prod_{l=1}^{i}(2^k-2^{i-l}) + a_i^{(n)}\cdot\prod_{l=1}^{i}(2^k-2^{i-l}) \\
&= \prod_{l=1}^{i}(2^k-2^{i-l})[a_i^{(n)}+a_{i-1}^{(n)}\cdot 2^{n-(i-1)}] = \prod_{l=0}^{i-1}(2^k-2^l)[a_i^{(n)}+2^{n-(i-1)}\cdot a_{i-1}^{(n)}].
\end{aligned}$$

On the other hand by George Landsberg [3] we have

$$\Gamma_i^{\left[\begin{smallmatrix} n\\ 1\end{smallmatrix}\right]\times k} = \prod_{l=0}^{i-1}\frac{(2^{n+1}-2^l)(2^k-2^l)}{(2^i-2^l)}. \tag{10.13}$$

Hence by (10.12) and (10.13) we have the formula

$$a_i^{(n)} + 2^{n-(i-1)}\cdot a_{i-1}^{(n)} = \prod_{l=0}^{i-1}\frac{2^{n+1}-2^l}{2^i-2^l}. \tag{10.14}$$

From (10.14) we deduce

(10.15)

$$\sum_{s=0}^{i-1}(-1)^s\cdot 2^{s(n-i)+\frac{s(s+1)}{2}}\left[a_{i-s}^{(n)}+2^{n-(i-(s+1))}\cdot a_{i-(s+1)}^{(n)}\right] = \sum_{s=0}^{i-1}\prod_{l=0}^{i-(s+1)}\frac{2^{n+1}-2^l}{2^{i-s}-2^l}(-1)^s\cdot 2^{s(n-i)+\frac{s(s+1)}{2}}.$$

Using (10.15) we get after some simplifications

$$a_i^{(n)} - (-1)^i a_0^{(n)} 2^{in-\frac{i(i-1)}{2}} = \sum_{s=0}^{i-1}\prod_{l=0}^{i-(s+1)}\frac{2^{n+1}-2^l}{2^{i-s}-2^l}(-1)^s\cdot 2^{s(n-i)+\frac{s(s+1)}{2}}.$$

$\square$

10.3. **Proof of Theorem 3.9.** Theorem 3.9 follows from Lemma 10.1 and Lemma 10.2.

10.4. **Proof of Corollary 3.10.** The assertions follows from Theorem 3.9 by some simple calculations.

10.5. **Proof of Theorem 3.11.** The proof of Theorem 3.11 is just a generalization of the proof of Lemma 6.9.

REFERENCES

[1] Daykin David E, *Distribution of Bordered Persymmetric Matrices in a finite field* J. reine angew. Math, **203**, 47-54 (1960).

[2] Hayes , D.R, *The expression of a polynomial as a sum of three irreducibles* Acta Arith. **11**, 461-488 (1966).

[3] Landsberg, G *Ueber eine Anzahlbestimmung und eine damit zusammenhangende Reihe*, J. reine angew. Math., **111**, 87-88 (1893).

Département de Mathématiques, Université de Brest, 29238 Brest cedex 3, France
E-mail address: `Jorgen.Cherly@univ-brest.fr`